




available at [www.sciencedirect.com](http://www.sciencedirect.com)



journal homepage: [www.elsevierhealth.com/journals/trst](http://www.elsevierhealth.com/journals/trst)



# Detection of HIV antigen and cDNA among antibody-negative blood samples in Nigeria

G.N. Odaibo<sup>a</sup>, A. Taiwo<sup>b</sup>, Y.A. Aken'Ova<sup>c</sup>, D.O. Olaleye<sup>a,\*</sup>

<sup>a</sup> Department of Virology, University College Hospital, College of Medicine, University of Ibadan, Ibadan, Nigeria

<sup>b</sup> Oyo State Blood Transfusion Unit, Total-Garden, Ibadan, Nigeria

<sup>c</sup> Department of Hematology, University College Hospital, College of Medicine, University of Ibadan, Ibadan, Nigeria

Received 3 July 2007; received in revised form 4 December 2007; accepted 4 December 2007

## KEYWORDS

HIV;  
Blood;  
Antigens;  
Antibodies;  
Blood transfusion;  
Diagnostic techniques  
and procedures

**Summary** In developing countries as many as 50% of patients for whom a transfusion is indicated are at risk of dying immediately if transfusion is withheld. It is therefore important that blood transfusion is made as safe as possible. This study was designed to assess the safety of blood transfusion in two large blood banks in Ibadan, Nigeria. Aliquots of 250 samples already screened and passed as negative for HIV-1 and -2 were collected from each of the blood banks. Samples were tested for the presence of HIV-1 antigen (ELAVIA Ag I) and the antigen-positive samples tested for the presence of specific HIV-1 antibodies by Western blot (BioRad, France). All antigen-positive samples were also subjected to PCR. HIV-1 antigen was detected in 6 (1.2%) of the 500 samples, of which 4 (0.8%) and 3 (0.6%) were Western blot-indeterminate and PCR-positive, respectively. Transfusion of HIV-contaminated blood may be contributing significantly to the spread of the virus in Nigeria. There is therefore an urgent need for an organized blood-banking system with facilities for more sensitive assays for the detection of HIV in blood to prevent transmission through transfusion.

© 2007 Royal Society of Tropical Medicine and Hygiene. Published by Elsevier Ltd. All rights reserved.

## 1. Introduction

The effectiveness of HIV intervention programmes, especially prevention of transmission through blood transfusion, depends largely on the accuracy of diagnosis of the infection. Blood transfusion is the most efficient mode of transmission of the virus. Reports have shown that over

90% of recipients of seropositive blood subsequently became infected (Simonsen et al., 1999). In developed countries, the rate of HIV transmission through transfusion has substantially reduced since 1985, when HIV antibody testing became available to screen blood and blood products for transfusion (Donegan et al., 2003). However, this is not the case in Africa, where blood transfusion is very common (Lee and Allain, 2004), even when it may not be medically indicated. According to Simonsen et al. (1999), injections are given when oral medicines would suffice and transfusion when more benign therapy should be preferred.

\* Corresponding author. Tel.: +234 803 324 1317;

fax: +234 2 241 3545.

E-mail address: [ibvirology@yahoo.com](mailto:ibvirology@yahoo.com) (D.O. Olaleye).

Although data on HIV transmission by blood transfusion in Africa are very scarce, several factors, including variability of the virus, facilities, cost and competence of laboratory personnel (Lee and Allain, 2004; Olaleye, 2006), are some of the problems of HIV diagnosis that may enhance this mode of transfusion in the region. Multiple subtypes and circulating recombinant forms are known to circulate in Nigeria (McCutchan et al., 1999; Ojesina et al., 2006), and the quality of HIV testing varies with the level of health care facilities, whether private, secondary or tertiary. In addition, recent reports have indicated that the use of contaminated syringes and needles as well as false-negative HIV-testing results may be contributing to the large number of infections in many developing countries (Odaibo et al., 1998, 2004; Simonsen et al., 1999). We assessed the safety of blood units for transfusion in two large blood banks in Nigeria.

## 2. Materials and methods

This study was carried out using samples from two large blood banks (blood bank I and blood bank II) in Ibadan, the capital of Oyo State, Nigeria, from March to October 2005. The two blood banks used the same procedures: a single ELISA for testing of blood units (GeneScreen Plus test kit; BioRad, Paris, France).

Aliquots of 250 whole blood samples (10 ml) from every fifth blood unit previously tested and recorded as HIV-1 and HIV-2-negative with the GeneScreen Plus ELISA were collected from each of the blood banks and transported in an isolated specimen box containing ice packs to the Department of Virology, University College Hospital, Ibadan, for further analysis. The samples were centrifuged and plasma separated in aliquots immediately in the laboratory on arrival. Both plasma and packed cells were stored at  $-20^{\circ}\text{C}$  until analysed.

### 2.1. HIV-1 antigen capture assay

All the plasma samples were tested for the presence of HIV-1 antigen using a commercial sandwich-type ELISA for p24 antigen detection (ELAVIA Ag I; Sanofi Diagnostics Pasteur, Marnes La Coquette, France). Briefly, each plasma sample was treated with a mild acid solution-ICS (ELAVIA Ag I immune complex dissociation reagent) to dissociate any antigen/antibody complexes. The treated samples were then transferred to reaction plates and the

test performed according to the manufacturer's instructions.

### 2.2. Western blot analysis

All HIV-1 antigen-positive plasma samples were further tested for the presence of specific HIV-1 antibodies using a commercially available Western blot kit, LAV BLOT 1 (BioRad, Paris, France). Samples and reagents were brought to room temperature and the test performed according to the manufacturer's instructions. Briefly, samples were added onto Western blot strips, and after washing, incubation with conjugate and colour development solution, the reaction was stopped by washing three times with distilled water. The presence of antibody to at least one envelope protein of HIV-1 was considered to be positive; the presence of any band that did not meet the criteria for positivity was taken as indeterminate; and no band at all was interpreted as negative.

### 2.3. DNA extraction and PCR analysis

DNA was extracted from the stored packed cells of the HIV-1 antigen-positive samples using the Qiagen Miniprep genomic DNA extraction kit (Qiagen Inc., Valencia, CA, USA) and then subjected to PCR. A set of primers [H1P4327 (TAAGACAGCAGTACAAATGGCAG) and H1P5155as (CTCTGTG-GCCCCTGGTCTTCT)]; supplied by the Institute of Tropical Medicine, Antwerp, Belgium was used to amplify a region of the HIV-1 pol gene. The PCR cycling protocol was 2 min at  $94^{\circ}\text{C}$  for denaturation, followed by 35 cycles of  $94^{\circ}\text{C}$  for 30 s,  $50^{\circ}\text{C}$  for 30 s and  $72^{\circ}\text{C}$  for 60 s and then  $72^{\circ}\text{C}$  for 7 min for final extension. The PCR product was subjected to gel electrophoresis in 2% agarose gel and viewed under UV light.

## 3. Results

In total, 500 HIV antibody-negative samples from the two blood banks (250 samples from each) were analysed (Table 1), and 6 (1.2%) were found to be positive for HIV-1 antigen by the antigen-capture ELISA. Further testing of the antigen-positive samples by Western blot showed that none was positive and 4 (0.8%) were indeterminate. However, 3 (0.6%) of the samples were found to be positive for HIV-1 antigen by PCR. The two Western blot-negative samples were also negative by PCR, whereas three of the Western blot-indeterminate samples were positive by PCR (Table 2).

**Table 1** Detection of HIV-1 antigen and cDNA (by PCR) in seronegative blood samples for transfusion in Ibadan, Nigeria

Site	No. samples (antibody-negative)	HIV-1 Ag No. (%) +ve	Analysis of HIV-1 Ag +ve samples			
			Western blot			PCR
			No. (%) +ve	No. (%) indeterminate	No. (%) -ve	No. (%) +ve
Blood bank I	250	4 (1.6)	0	3 (1.2)	1 (0.4)	2 (0.8)
Blood bank II	250	2 (0.8)	0	1 (0.4)	1 (0.4)	1 (0.4)
Total	500	6 (1.2)	0	4 (0.8)	2 (0.4)	3 (0.6)

**Table 2** Western blot and PCR reactivity pattern of HIV-1 antigen-positive samples

Sample ID no.	HIV-1 Ag	Western blot	PCR
001	+ve	Indeterminate	+ve
002	+ve	Indeterminate	–ve
003	+ve	–ve	–ve
004	+ve	Indeterminate	+ve
005	+ve	–ve	–ve
006	+ve	Indeterminate	+ve

#### 4. Discussion

HIV infection resulting from blood transfusion of infected blood has been documented since the first case report in 1981 (Curran et al., 1985). The results of this study show that HIV-1 antigens were detected in 6 (1.2%) of 500 antibody-negative blood units, which was further confirmed by PCR in 3 (0.6%) of the samples. Similar observations have been reported in other parts of the world (Stramer et al., 2004). Failure to detect HIV antibodies in blood units donated by HIV-infected people who have not seroconverted has been documented (Nelson et al., 1999; Sitas et al., 1994). Significant changes in the antigenic structure of HIV may affect the sensitivity of serological tests, a likely scenario when circulation of multiple subtypes of the virus takes place, as is the case in Nigeria (McCutchan et al., 1999; Odaibo et al., 2006; Ojesina et al., 2006; Peeters et al., 2003).

The rate of unidentified HIV-1-positive blood units found in this study is considered to be high compared with reported cases in developed countries. For instance, only 6 of 8710 (0.07%) Western blot-indeterminate samples and 0 of 7562 Western blot-negative samples were confirmed positive for HIV RNA in the United States (Stramer, 2004). The findings in this study indicate that transfusion of HIV-contaminated blood may be contributing to the spread of the virus in Nigeria and that there is a need for urgent and necessary action by the relevant health authorities to prevent transfusion-associated transmission. The inclusion of P24 antigen testing and the nucleic acid amplification test, which detects HIV-1 RNA in minipools, has been shown to further reduce the risk of transfusing HIV-contaminated blood in the United States (Stramer, 2004).

It is interesting that none of the six antigen-positive samples was confirmed by Western blot, although four were indeterminate. This may indicate that the donors were recently infected with low levels of antibody or that they were infected with virus variants of the epitopes used for the antigen base of the assay kit. The reduced sensitivity of commercial diagnostic assays is particularly problematic in regions where multiple HIV-1 non-B subtypes and recombinants are predominant, as in Nigeria (Horal et al., 1991; Peeters et al., 2003). Thus there is a need for continuous monitoring of the performance of diagnostic assays, especially with a view to incorporating the structures of new subtypes/variants in the antigenic base. Although the indeterminate results may be due to the presence of other related, but different, retroviruses in the area, amplification of the pol gene fragment of HIV from three of the four

indeterminate samples confirmed the presence of the virus in some of the blood units.

The inability to amplify HIV cDNA from three (50%) of the antigen-positive samples may be due to primer mismatch caused by variation in the sequence of the virus. Apart from the cost and lack of facilities, sequence diversity of HIV has been a constant challenge for the application of PCR in African countries, where multiple genetic types and subtypes co-circulate (Fonjongo et al., 2000; Peeters et al., 2003). Nevertheless, PCR is a potentially valuable tool to diagnose HIV in donated blood and thus reduce the risk of transmission through blood transfusion.

It is also worthy of note that the blood banks where samples were collected for this study are in an urban setting where facilities and manpower can be considered close to adequate. The role of blood transfusion in semi-urban and rural communities, where there are little or no facilities available for pre-transfusion HIV screening, is likely to be much more significant. Thus it is necessary to put in place centralized blood-banking systems by state or region, where facilities for more sensitive assays for detection of HIV antibody/antigen in blood units for transfusion can be provided and to ensure proper monitoring through links or networking with HIV reference laboratories in Nigeria as well as other African countries.

#### Authors contributions

GNO and DOO designed the study protocol, carried out antigen detection and molecular analysis and interpreted the data; AT and YAA carried out sample collection and screening of blood units at the respective blood banks; GNO drafted the manuscript and DOO revised it. All authors read and approved the final manuscript. DOO is guarantor of the paper.

**Acknowledgements:** We appreciate the support of the other staff of the Virology Department, University College Hospital, Ibadan, Nigeria, and the two blood banks for their assistance during the course of the study. The primers used for the study were kindly provided by Dr W. Jassens' team at the Department of Microbiology, Institute of Tropical Medicine, Antwerp, Belgium.

**Funding:** Funding for this study was provided by the HIV/Viral Hepatitis Project of the Department of Virology, College of Medicine, University of Ibadan, Ibadan, Nigeria.

**Conflicts of interest:** None declared.

**Ethical approval:** University of Ibadan/University College Hospital Ethical Review Committee, Ibadan, Nigeria.

#### References

- Curran, J.W., Lawrence, D.N., Jaffe, H., 1985. AIDS associated with transfusion. *N. Engl. J. Med.* 310, 6–75.
- Donegan, E., Moncada, J., Chernesky, M., Schachter, J., 2003. Sexually transmitted infection/HIV laboratory testing in resource-limited countries: steps to improve use effective testing and to ensure diagnostic capacity. *Sex. Transm. Dis.* 30, 905–908.

- Fonjungo, P.N., Mondji, E.N., Torimiro, J.N., Eno, L.T., Nkengkong, J.N., Gao, F., Rayfield, M., Folks, T.M., Pieniazek, D., Lal, R.B., 2000. Presence of diverse human immunodeficiency virus type 1 viral variants in Cameroon. *AIDS Res. Hum. Retroviruses* 16, 1319–1324.
- Horat, P., Svennerholm, B., Jeansson, S., Rymo, L., Hall, W.W., Vahlne, A., 1991. Continuous epitopes of human immunodeficiency virus type 1 (HIV-1) transmembrane glycoprotein and reactivity of human sera to synthetic peptides representing various HIV-1 isolates. *J. Virol.* 65, 2718–2723.
- Lee, H.H., Allain, J.P., 2004. Improving blood safety in resource-poor settings. *Vox Sang.* 87, 176–179.
- McCutchan, F.E., Carr, J.K., Bajani, M., Sanders-Buell, E., Harry, T.O., Stoeckli, T.C., Robbins, K.E., Gashau, W., Nasidi, A., Janssens, W., Kalish, M.L., 1999. Subtype G and multiple forms of A/G intersubtype recombinant human immunodeficiency virus type 1 in Nigeria. *Virology* 254, 226–234.
- Nelson, K.E., Rungtongthani, K., Margolick, J., Suriyanon, V., Niyomthai, S., Boer, M.A., Kawichai, S., Robison, V., Celentano, D.D., Nagachinta, T., Duerr, A., 1999. High rates of transmission of subtype E human immunodeficiency virus type 1 among heterosexual couples in northern Thailand: role of sexually transmitted disease and immune compromise. *J. Infect. Dis.* 180, 337–343.
- Odaibo, G.N., Olaleye, O.D., Tomori, O., 1998. Human immunodeficiency virus types 1 and 2 infection in some rural areas of Nigeria. *Rom. J. Virol.* 49, 89–95.
- Odaibo, G.N., Bamgbose, G., Jegede, A.S., Sankale, J.-L., Omotade, O.O., Olaleye, D.O., Kanki, P., 2004. Demographic and Laboratory Evidence of Non Sexual Transmission of HIV in Nigeria, in: *Published Proceedings of the XV International AIDS Conference E710L7663*, Medimond S.r.l., Bologna, Italy, pp. 25–30.
- Odaibo, G.N., Olaleye, D.O., Heyndrickx, L., Vereecken, K., Houwer, K., Janssens, W., 2006. Mother-to-child transmission of different HIV-1 subtypes among ARV naïve infected pregnant women in Nigeria. *Rev. Inst. Med. Trop. Sao Paulo* 8, 77–80.
- Ojesina, A.L., Sankale, J.L., Odaibo, G., Langevin, S., Meloni, S.T., Sarr, A.D., Olaleye, D., Kanki, P.J., 2006. Subtype-specific patterns in HIV type 1 reverse transcriptase and protease in Oyo State, Nigeria: implications for drug resistance and host response. *AIDS Res. Hum. Retroviruses* 22, 770–779.
- Olaleye, D.O., 2006. National HIV Testing Algorithm. Federal Government of Nigeria, Nigeria.
- Peeters, M., Toure-Kane, C., Nkengasong, J.N., 2003. Genetic diversity of HIV in Africa: impact on diagnosis, treatment, vaccine development and trials. *AIDS* 17, 2547–2560.
- Simonsen, L., Kane, A., Lloyd, J., Zaffran, M., Kane, M., 1999. Unsafe injections in developing world and transmission of blood-borne pathogens. *Bull. World Health Organ.* 77, 789–800.
- Sitas, F., Fleming, A.F., Morris, J., 1994. Residual risk of transmission of HIV through blood transfusion in South Africa. *South Afr. Med. J.* 84, 142–144.
- Stramer, S.L., 2004. Viral diagnostics in the arena of blood donor screening. *Vox Sang.* 87 (Suppl. 2), 180–183.
- Stramer, S.L., Glynn, S.A., Kleinman, S.H., Strong, D.M., Caglioti, S., Wright, D.J., Dodd, R.Y., Busch, M.P., 2004. National Heart, Lung, and Blood Institute Nucleic Acid Test Study Group Detection of HIV-1 and HCV infections among antibody-negative blood donors by nucleic acid-amplification testing. *N. Engl. J. Med.* 351, 760–768.

UNIVERSITY OF IBADAN LIBRARY