

# NIGERIAN JOURNAL OF ANIMAL PRODUCTION



*Published by*  
**THE NIGERIAN SOCIETY FOR ANIMAL PRODUCTION**

*with the support of*



**VOLUME 39**

**2012 NUMBER 2**

# NIGERIAN JOURNAL OF ANIMAL PRODUCTION

Editor-In-Chief

Dr. O. A. Adeyemi

Deputy Editor-In-Chief

Dr. O. J. Babayemi

## OFFICERS OF THE NIGERIAN SOCIETY FOR ANIMAL PRODUCTION 2012-2014

President

Prof. C.F.I. Onwuka

Vice President

Dr. A. O. Ani

Secretary

Dr. F. O. Abeke

Assistant Secretary

Dr. (Mrs.) L.A.F. Akinola

Editor-In-Chief

Dr. O. A. Adeyemi

Deputy Editor-In-Chief

Dr. O. J. Babayemi

Treasurer

Dr. S. B. Abdu

Public Secretary

Dr. A.O. Abu

Ex-Officio

Prof. Udo Herbert

### Editorial Board

Prof. Udo Herbert Dr. O. A. Adeyemi Dr. O. J. Babayemi Dr. A. A. Taiwo

The Nigerian Society for Animal Production was inaugurated in March 1973 during the first International Symposium on Animal production in the Tropics at the University of Ibadan, Ibadan, Nigeria. This Society is responsible for the publication of the Nigerian Journal of Animal Production which commenced the publication in 1974. The journal publishes two issues in a year. This present issue features the biography of one eminent personality as mark of honour to him.

### Membership

Membership of the Society is open to all that are engaged or interested in any aspect of Animal Production including Fisheries and Wildlife Management from all over the world. Interested individuals organizations or institutions should apply for membership to the Secretary of the Society.

### Subscription and Advertising:

The current annual subscription is N1,500 for non-members and N2000 for organization and institutions within the country. It costs one hundred and twenty eight US Dollars (US \$128) for subscription outside Nigeria. Correspondence concerning subscriptions and purchase of single or back copies of the journal should be addressed to the Deputy Editor-in Chief, Nigerian Journal of Animal Production Dr. O. J. Babayemi, Department of Animal Science University of Ibadan, Ibadan, Nigeria. E-mail: oj.babayemi@mail.ui.edu.ng.

### Manuscripts

All manuscripts and correspondence concerning the journal should be sent to Editor-in-Chief, Dr O.A. Adeyemi, Department of Animal Production and Health, Federal University of Agriculture, P.M.B. 2240, Abeokuta. Email: olajideadeyemi@yahoo.com. Manuscripts are invited from all over the world as long as they are original research reports on any aspect of Animal Production and Health. All authors' name in the references must be in bold lower case letters except the first letter of each surname. The surname must be followed by the initials in all cases irrespective of the number of authors. Tables and legends for figures must be typed on separate sheets. Tables must be numbered, title written in lower case letter (except the first letter) and left justified. Horizontal and vertical lines within the tables are not acceptable. Submissions on computer disc(CD) are also acceptable.

- Prof. CEL. Onwuka
- Dr. A. O. Adin
- Dr. F. O. Aboki
- Dr. (Mrs) J. A. F. Akinola
- Dr. O. A. Adesomi
- Dr. O. J. Babayemi
- Dr. S. B. Abdu
- Dr. A. O. Abdu
- Prof. Udo Herbert

# NIGERIAN JOURNAL OF ANIMAL PRODUCTION

- President
- Vice President
- Secretary
- Assistant Secretary
- Editor-in-Chief
- Deputy Editor-in-Chief
- Treasurer
- Public Secretary
- Ex-Officio

ISBN 0331-2064

Prof. Udo Herbert Dr. O. J. Babayemi Dr. A. A. Taiwo

Typeset and formatted by

**Azol Computers**

Department of Animal Production and Health

University of Agriculture, Abeokuta

08036250504

The Nigerian Society for Animal Production is a non-profit organization established in 1975 during the first international symposium on animal production in Lagos, Nigeria. The Society is responsible for the publication of the Nigerian Journal of Animal Production since 1974. The journal publishes research papers on all aspects of animal production in Nigeria and other developing countries. The journal is published quarterly as a means of disseminating information to the members of the Society.

Printed by

**Lynson Prints**

No, 31 Sokenu Rd.

Oke-Ijeun, Abeokuta

08033928540, 08175608494

Published by

## THE NIGERIAN SOCIETY FOR ANIMAL PRODUCTION

Abstracted in (I) CABS

(ii) African Journal Online (ajol)

<http://www.inasp.info/ajol>

**AKNOWLEDGEMENT OF THE REVIEWERS OF ARTICLES IN  
VOLUME 39, (NUMBER 2) 2012 OF THE NIGERIAN JOURNAL OF  
ANIMAL PRODUCTION**

The Editorial Board wishes to express appreciations and thanks to the following reviewers who have helped in the assessment of articles published in Volume 39, Number 2, 2012 of the Nigerian Journal of Animal Production

Adeniji, A. A. (Dr.)	Iyeghe-Erakpotobor, G. T. (Prof. Mrs.)
Adeyemi, O. A. (Dr.)	James, I. J. (Dr.)
Agwunobi, L. N. (Prof.)	Odugbemi, T. (Dr.)
Akinola, O. S. (Dr.)	Ogunbosoye, D. (Dr.)
Ayasan, T. (Dr.)	Okagbare, G. O. (Prof.)
Babayemi, O. J. (Dr.)	Olugbemi, T. A. (Dr.)
Bamiklolle, M. A. (Prof.)	Oluremi, O. I. A. (Prof.)
Bemji, M. (Dr.)	Omoyakhi, Moses (Dr.)
Ebenso, Ime (Dr.)	Oni, Jentola (Dr.)
Esonu, B. O. (Prof.)	Orheruata, A. M. (Prof.)
Etuk, E. (Dr.)	Shittu, A. M. (Dr.)
Ezeokeke, C. T. (Dr.)	Sogunle, O. M. (Dr.)
Idowu, F. (Dr.)	Talabi, A. O. (Dr.)

TABLE OF CONTENTS

CONTENTS	PAGES
<b>A brief biography</b> <b>Professor Jerome O. Gefu</b> <b>Animal Breeding and Genetics</b>	<b>1</b>
Effects of genotype and season on the productive performance of commercial egg-type chickens in the derived savanna zone in Nigeria <b>A.O. Oguntunji and A.E. Salako</b>	<b>4</b>
Biometric measurements of body and internal organ traits of indigenous chickens on free range in South-Eastern Nigeria <b>V. M. O. Okoro, G. E. Ejiogu, I. C. Okoli, G. A. Anyanwu, C. T. Ezeokeke and C. L. Okoro</b>	<b>14</b>
Udder and linear body measurement in ewes and does in relation to performance of their offspring under the traditional system of management <b>O.O. Adewumi, O. Banjo, A.A. Adegboyega and O.A. Noiki</b>	<b>24</b>
Heterotic effect on body weight and morphometric traits of crossbred buck kids of Red Sokoto and West African Dwarf goats <b>E.N. Nwachukwu and G.U. Okoji</b>	<b>35</b>
Performance of pure and crossbred progenies of Red Sokoto and West African Dwarf goats in the Rainforest Zone of South Eastern Nigeria <b>E. N. Nwachukwu, K. U. Amaefule, F. O. Ahamefule, S.C. Akomas, T. U. Nwabueze, U. A. U. Onyebinama and O. O. Ekumankama</b>	<b>41</b>
<b>Animal Physiology</b>	
Composition and specific gravity of milk of West African Dwarf sheep as affected by stage of lactation and parity <b>T.J. Williams, I.J. James, M.R. Abdulateef, L.O. Onabegun, S.O. Jinadu, Y.O. Falade, F.T. Solola, O.O. Adewumi and O.E. Oke</b>	<b>49</b>

## Non-Ruminant Nutrition, Production and Management

- Comparative chemical composition of 24-hour fermented sweet orange fruit (*Citrus sinensis*) peel meal and maize and effect on performance response of starting pullet chicks  
**B.O. Oyewole, O.I.A. Oluremi, S.O. Aribido and J.A. Ayoade** 57
- Performance of grower rabbits fed *Gmelina arborea* leaf meal  
**A. A. Adeniji, E. Gana, O. A. Olafadehan, I.C. Chibougwu and M. I. Saria** 66
- Performance, carcass and blood indices of broiler chickens fed graded levels of sundried raw wild cocoyam (*Colocasia esculenta* (L.) Schott) corms as partial substitutes for maize  
**R. Olajide** 71
- Effect of  $\beta$ -mannanase supplementation and feed presentation on carcass characteristics and macro-mineral digestibility of growing rabbits fed palm kernel based diets  
**O. A. Abu and D. A. Tanimowo** 83
- Cost benefit analysis of level of Ziziphus (*Ziziphus mauritiana*) leaf meal inclusion in the diet of growing rabbits  
**S.B. Abdu, M.R. Hassan, H.Y. Adamu, S.M. Yashim, M. Abdulrashid, S. Duru and G.E. Jokthan** 90
- Influence of feed quantity offered on growth performance, carcass yield, organs weight and back-fat composition of finishing pigs  
**C.P. Njoku, A.B.J. Aina, O.M. Sogunle, O.A. Adeyemi, and O.O. Oduguwa** 96
- Physiological evaluation of broiler finishers fed fermented bovine blood and rumen digesta meal  
**B.O. Esonu, J.C. Azubuike, A.B.I. Udedibie, M. N. Opara and T. C. Iwuji** 107
- Effects of enzyme supplementation of raw bambara nut (*Vigna subterranea* (L) verdc) waste diets on carcass and organ weights of broiler finishers  
**A. O. Ani and O.D. Omeje** 114
- Effects of diets containing alkali-treated soybeans on performance traits, nutrient digestibility and cost benefits of broiler chickens  
**M. M. Ari, B.A. Ayanwale, T .Z. Adama and E.A. Olatunji** 125

## Ruminant Nutrition, Production and Management

Unethical practices relating to cattle freedom, care and control among the market handlers in Ibadan metropolis, Nigeria

**P. L. Akinyemi, O. J. Babayemi, C. Agulana and M. K.**

**Bamikole**

135

*In - vitro* fermentation and chemical constituents of urea-molasses feed - blocks made with different binders for ruminants

**O. A. Isah**

142

## Fisheries and Aquaculture Management

Water quality and fish diversity of Federal University of Agriculture Abeokuta reservoir

**F.I. Adeosun, W.O. Abdul, A.A. Akinyemi, A.O. Agbon and**

**O.C. Odebiyi**

150

Evaluation of fermented pineapple (*Ananas comosus*) peel meal as a substitute for maize meal in the diet of Nile tilapia, *Oreochromis niloticus*

**S.O. Obasa, W.O. Alegbeleye, F.O.A. George, W.A. Abdul, N.A. Bamidele,**

**M.T. Okon and W.I. Olaleye**

156

## Animal Health

Evaluation of the toxic effect of aqueous extract of the bark of *Morinda morindoides* root in male wistar rats

**J.O. Olukunle, O.L. Ajayi, O.T. Adenubi, E. B. Jacobs, B.S. Okediran**

**and O. A. Akinkuotu**

163

Effects of *Lippia multiflora* leaf extract on the semen of wistar rat

**J.O. Olukunle, E. S. Ajibola, K.T. Biobaku, B.S. Okediran and**

**O. A. Akinkuotu**

172

## Micro Livestock Production

Influence of different management systems on gut microbes and chemical constituents of giant land snail (*Archachatina marginata*)

**K.O. Ademolu, A. B. Idowu, O.A. Jayeola, I. Osunsina, G..A. Dedeke, F. Oluwafemi,**

**and E. Ibie**

179

## **Animal and Livestock Products / Processing**

Quality characteristics of chevon obtained from goat fed melon (*Colocynthis citrillus*) husk and palm (*Elaeis guineensis*) oil slurry

**K.A. Sanwo, S.O. Iposu1, J.A. Adegbite, S.S. Abiola, A.O. Fanimo and O.B. Oyewole**

187

## **Livestock Economics / Extension**

Technical efficiency of local dairy milk processing in Kwara State, North Central, Nigeria

**E. O. Olorunsanya and A. M. Daramola**

194

Consumers preference for rabbit meat as a source of animal protein in selected Local Government Areas in Rivers State

**L.A.F. Akinola and M.O. Ironkwe**

203

## **Feed Resources**

Visual assessment, proximate composition and cost analysis of three differently processed discarded vegetable-bovine blood-rumen content mixtures

**D.A. Ekunseitan, O.O. Balogun, D. Erubetina, S.S. Abiola, O.M. Sogunle, I.M.Ogunade, L.T. Egbeyale, A.A. Ayoola, O.F. Akinola, I. B. Allison and S.O. Osho**

211

Effects of raw and toasted *Daniellia oliveri* (Rolfe) seed meal on broiler chicken performance

**C.O. Obun, and O. A. Adeyemi**

218



**Effect of  $\beta$ -mannanase supplementation and feed presentation on carcass characteristics and macro-mineral digestibility of growing rabbits fed palm kernel based diets**

O. A. Abu and D. A. Tanimowo

Agricultural Biochemistry and Nutrition Unit, Department of Animal Science, University of Ibadan, Ibadan, Nigeria

Corresponding author's E-mail: [oa.abu@mail.edu.ng](mailto:oa.abu@mail.edu.ng); [ohiahmed@yahoo.com](mailto:ohiahmed@yahoo.com)  
Phone No +2348058009872**Abstract**

The influence of  $\beta$ -mannanase supplementation and feed presentation of palm kernel cake (PKC) based diets were assessed on grower rabbits. Twenty crossbred grower rabbits ( $757.5 \pm 65$ g) of mixed sex aged between eight to ten weeks old were used during the trial in 2 X 2 factorial design conducted in a completely randomized experimental design giving four treatment interactions with five replicates of one rabbit per replicate. The treatments were; T1- diet in mash form without enzyme; T2- diet in pellet form without enzyme; T3- diet in mash form with  $\beta$ -mannanase (0.5g/kg) and T4- diet in pellet form with mannanase (0.5g/kg). The trial lasted for six weeks. The animals were weighed individually at the beginning of the study and weekly thereafter. At the end of the study period the faecal samples were collected and animals were slaughtered and measurements taken from the carcasses. The carcass yield showed no interaction, though full GIT weights (g) were significantly influenced (T1-23.04, T2-17.67, T3-19.99 and T4- 18.74). Enzyme inclusion and feed form elicited significant effects on carcass characteristics, with varying improvements observed. Enzyme inclusion on the mash diet gave lower organ weights compared to other diets. Feed form and enzyme inclusion influenced lengths of the segments of the GIT, with rabbit on Diet 3 having the longest length. Pelleting influenced ileum lengths positively. Rabbits fed pelleted or mash diets whether supplemented or non-supplemented with  $\beta$ -mannanase digested dry matter, calcium, phosphorus and magnesium in the same level. In conclusion, supplementation of diets with  $\beta$ -mannanase and pelleting of diets improved phosphorus digestibility in palm kernel cake based diets. In addition, rabbits feed mash and pelleted rations with addition of  $\beta$ -mannanase had improved loins as opposed to those fed mash and non-supplemented rabbits while other carcass and visceral organ characteristics were not significantly affected.

**Keywords:**  $\beta$ -mannanase, palm kernel cake, feed presentation, macro-minerals, carcass characteristics, rabbits

**Introduction**

A stiff competition exists between humans and the livestock industry for cereal grains, and secondly, the low production of grains results in its scarcity (Oluwafemi, 2009) and hence more reliance of livestock on agro-industrial by-products. One of such alternatives is Palm Kernel Cake (PKC). The Palm Kernel Cake (PKC) is a potential feedstuff obtained from the oil-palm processing industry. Palm kernel cake is a by-product of the extraction of oil from the

palm kernels of the oil palm (*Elaeis guineensis*). Oil from palm kernel can be extracted either manually or by appropriate organic solvents. Palm kernel cake obtained during manual extraction of palm kernel oil is of higher value because of the residual oil left unlike the solvent extracted that contains little or no residual oil. With a chemical composition close to rice bran and wheat bran, it has a potential for feeds. Its use has however been limited by its low protein, highly fibrous and gritty nature (Longe and Fagbenro-Bryon, 1990) and

presence of non-starch polysaccharides (NSPs). These NSPs impair the digestibility and utilization of nutrients present in the plant feedstuff (Han, 1996). However, recent advances in biotechnology have opened opportunities for improvement of palm kernel cake for livestock utilization (Ng and Chong, 2002). The  $\beta$ -mannan is the main NSP found in palm kernel cake (Sundu *et al.*, 2006) which is reported to be as high as 30% in the palm kernel meal (ChemGen Corp).

In relatively recent years there has been a renewed interest in the use of exogenous enzymes (feed grade enzymes) or solid state fermentation (Mosobalaje *et al.*, 2012) to enhance the nutritive value of a diet, through their action on the utilization of nutrients, as well as improved health of a flock (Sarvestani *et al.*, 2006).

Earlier works have shown that usage of feed in the form of pellets have physical and nutritional benefits (Behnke, 1994, 2001, Salari *et al.*, 2006). The physical benefits include improved ease of handling, reduced ingredient segregation, less feed wastage and increased bulk density. Falk, 1985, reported some of the nutritional benefits of feeding pellets. The improvements in performance of rabbits fed pelleted diets have been attributed to decreased feed wastage, reduced selective feeding, less time and energy expended for prehension, destruction of pathogens, thermal modification of starch and improvement in palatability (Behnke, 1994) and not as a result of fat deposition (Hathorn and Mowlem, 1980). Increase in cost of pelleting must be off set by improved performance for profitability (Ziggers, 2004). To date, studies on enzyme inclusion in rabbit diets in Nigeria have been inconclusive, while studies on enzyme inclusion in pelleted diets is lacking. The purpose of this study was to determine the

influence of  $\beta$ -mannanase and feed form (pellet and mash) on the performance of rabbits on a palm kernel cake based diet.

### Materials and methods

Four isonitrogenous (17%) and isoenergetic (10.0 MJ<sup>l</sup>) practical diets were formulated (Table 1). All the diets contained 30% PKM as energy source. Two of the diets were in mash form while the other two were pelleted. Diet 3 and 4 were supplemented with  $\beta$ -mannanase (ChemGen Corp.) at 0.05% inclusion level.  $\beta$ -mannanase was added along with the micronutrients to both diets during the milling process

### Experimental procedure

Twenty grower rabbits with a mean initial weight of 757.5±65g were randomly allocated to the four treatments with five (5) replicates per treatment, in a 2 x 2 factorial treatment design in a completely randomized design (CRD). At the start of the experiment, rabbits were weighed and allocated to individual cells of 40cm x 45cm x 60cm (L x B x H). The hutches were placed indoors with dwarf walls with concrete impervious floor. The rabbits were fed in the mornings and evenings. Water was given *ad-libitum*. Leftovers were removed and weighed daily. The rabbits were weighed weekly until the last week of the experimental period. The feeding trial lasted for six weeks.

### Evaluation and analysis

At the beginning of the 6<sup>th</sup> week, two rabbits per replicate for each treatment were put in metabolic cages for 4 days for digestibility trial. Feed intake measured and fecal output was collected for each of the 4 days, weighed and recorded for each replicate, from which an aliquot (10%) taken and oven dried. Calcium, phosphorus and magnesium were measured by dissolving the samples in nitric acid and hydrochloric

Table 1. Composition of the experimental diets

Ingredients	Diet 1	Diet 2	Diet 3	Diet 4
	(Mash)	(Pellet)	(Mash) + enzyme	(Pellet) + enzyme
Palm kernel cake	30.00	30.00	30.00	30.00
Maize	18.00	18.00	18.00	18.00
Groundnut cake	12.00	12.00	12.00	12.00
Wheat bran	11.00	11.00	11.00	11.00
Corn bran	10.00	10.00	10.00	10.00
Cowpea husk	7.00	7.00	7.00	7.00
Rice husk	7.00	7.00	7.00	7.00
Cassava starch	2.00	2.00	2.00	2.00
Limestone	1.50	1.50	1.50	1.50
Dicalcium phosphate	0.50	0.50	0.50	0.50
Table salt	0.25	0.25	0.25	0.25
DL-Methionine	0.25	0.25	0.25	0.25
L-Lysine	0.20	0.20	0.15	0.15
Hemicell (Mannanase)	0.00	0.00	0.05	0.05
Premix	0.25	0.25	0.25	0.25
Mycofix	0.05	0.05	0.05	0.05
<b>Total</b>	<b>100.00</b>	<b>100.00</b>	<b>100.00</b>	<b>100.00</b>
<b>Calculated chemical composition</b>				
Crude protein (%)	17.12	17.12	17.12	17.12
Crude fiber (%)	11.53	11.53	11.53	11.53
Digestible energy (MJ/kg)	9.97	9.97	9.97	9.97

## \*Composition of Premix

Vit A- 3,000,000iu/kg, Vit E-3000iu/kg, Vit B<sub>1</sub>-200mg/kg, Vit B<sub>6</sub>-80mg/kg, Niacin-6000mg/kg, Molybdenum-200mg/kg, Iron-16,000mg/kg, Cobalt-100mg/kg, Zinc-20000mg/kg, Copper-2400mg/kg, Vit D<sub>3</sub>-600000iu/kg, Vit K<sub>3</sub>-800mg/kg, Vit B<sub>12</sub>-2000mg/kg, Biotin-20000mcg/kg, Iodine-200mg/kg, Selenium-40mg/kg, Manganese-40000mg/kg, Pantothenic acid-2400mg/kg

acid and diluting with de-ionized water to 100ml. The resultant extracts analyzed for calcium and magnesium using an atomic spectrophotometer and phosphorus using an autoanalyzer. At the end of the 6<sup>th</sup> week, all the rabbits were slaughtered, flayed and the carcasses divided into various parts and evaluated. Morphometric measurements were carried out on the gastro-intestinal tract (GIT) of the rabbits to determine the effects of the treatments on the each section.

**Statistical analysis**

The data collected was subject to analysis of variance, using the general linear model (GLM) procedure of SPSS (2001) and where significant was subject to the Duncan's Multiple Range Test (Duncan,

1955).

**Results**

Presented in Table 2 are the values for carcass evaluation. Diet 3 produced the highest values for dressing percentage, head and shoulder weights ( $P > 0.05$ ). Diet 2 however gave a higher value for pelt, rack ( $P < 0.05$ ) and thighs ( $P > 0.05$ ). Loin weights revealed that Diet 4 gave a significantly higher weight than Diets 2 and 3, Diet 3 was also significantly different from Diet 1. No significant differences were observed among the treatments for all the visceral organ weights (Table 3). Diet 1 gave the highest values for liver, abdominal fat, kidney and spleen while the lowest

Table 2. Carcass characteristics as influenced by feed form and  $\beta$ -mannanase supplementation on grower rabbits

Parameters (%)	Diet 1	Diet 2	Diet 3	Diet 4	SEM	Significance
Dressing out	48.22	50.78	52.02	51.09	1.86	ns
Pelt	9.17 <sup>ab</sup>	10.42 <sup>a</sup>	8.54 <sup>b</sup>	9.11 <sup>ab</sup>	0.60	*
Head	9.11	9.38	9.62	9.28	0.35	ns
Shoulders	7.36	7.26	10.72	8.59	1.31	ns
Thighs	17.56	17.59	16.84	16.82	0.75	ns
Rack	10.58 <sup>ab</sup>	10.95 <sup>a</sup>	8.20 <sup>b</sup>	9.11 <sup>b</sup>	0.59	**
Loins	12.43 <sup>c</sup>	14.61 <sup>bc</sup>	15.74 <sup>ab</sup>	16.39 <sup>a</sup>	0.92	*

a, b – means in the same row with different alphabets are significantly different.

(%) Percentages are based on live weight

values for liver, abdominal fat, kidney, spleen and lungs were observed on Diet 3. The highest values for the heart and lungs were on Diet 2.

Values for GIT evaluation showed significant differences in total gastrointestinal tract length (Table 5). There was however difference in the duodenum, jejunum and caecum lengths. Diet 4 had the highest value for stomach length ( $P < 0.05$ ). Ileum length was also influenced by the association between feed form and mannanase inclusion with Diet 2 having the

highest value ( $P < 0.05$ ), the same trend was observed for large intestine and colon lengths ( $P < 0.05$ ). There was however no difference in the caecum lengths of the rabbits, though Diet 1 was observed to have the highest value (Table 4).

Macro mineral digestibility showed no difference in the values for calcium and magnesium, though Diet 4 had numerically higher values for dry matter and calcium (Table 5). Digestibility of phosphorus ( $P < 0.05$ ) and magnesium ( $P > 0.05$ ) was highest on Diet 3. Diet 2 was observed to give the lowest digestibility for all the

Table 3. Effect of feed form and Mannanase on visceral organ weights of grower rabbits

Parameters (%)	Diet 1	Diet 2	Diet 3	Diet 4	SEM	Significance
Heart	0.29	0.33	0.32	0.30	0.02	ns
Liver	3.17	2.75	2.72	2.78	0.21	ns
Lungs	0.56	0.60	0.48	0.60	0.08	ns
Abdominal fat	1.03	0.81	0.71	0.84	0.21	ns
Kidney	0.60	0.57	0.51	0.57	0.04	ns
Spleen	0.06	0.05	0.04	0.04	0.01	ns

(%); Percentages based on live weight

Table 4. Influence of feed form and mannanase supplementation on GIT parameters of rabbits

Parameters	Diet 1	Diet 2	Diet 3	Diet 4	SEM	Significance
GIT Length (m)	3.46 <sup>b</sup>	3.86 <sup>ab</sup>	4.16 <sup>a</sup>	3.89 <sup>a</sup>	0.17	*
Stomach (cm)	15.60 <sup>c</sup>	16.44 <sup>bc</sup>	18.62 <sup>b</sup>	23.16 <sup>a</sup>	1.08	**
Duodenum (cm)	42.60	48.40	58.10	42.44	4.01	ns
Jejunum (cm)	71.00	66.54	75.84	68.60	7.85	ns
Ileum (cm)	88.60 <sup>b</sup>	117.0 <sup>a</sup>	87.38 <sup>b</sup>	82.40 <sup>b</sup>	9.32	*
Large intestine (cm)	28.64 <sup>b</sup>	37.90 <sup>a</sup>	31.50 <sup>ab</sup>	31.40 <sup>ab</sup>	2.62	*
Caecum (cm)	49.90	47.32	49.36	46.98	4.07	ns

a, b, c – means in the same column with different alphabets are significantly different

GIT – gastro-intestinal tract length

**Table 5. Influence of feed form and mannanase supplementation on dry matter, calcium, phosphorus and magnesium digestibility in grower rabbits**

Parameters	Diet 1	Diet 2	Diet 3	Diet 4	SEM	Significance
Dry matter	0.62	0.58	0.64	0.65	0.037	ns
Calcium	0.67	0.64	0.68	0.71	0.032	ns
Phosphorus	0.53 <sup>ab</sup>	0.41 <sup>b</sup>	0.65 <sup>a</sup>	0.61 <sup>a</sup>	0.046	*
Magnesium	0.41	0.40	0.52	0.49	0.055	ns

a, b, c – means in the same column with different subscripts are significantly different

metabolites.

### Discussion

The non-significant effects on most of the visceral organs and carcass characteristics of simple stomach animals fed enzyme-supplemented rations have been reported (Garcia *et al.*, 2003; Iyayi and Davies, 2005). These findings were contrary to Olayemi *et al.* (2006) who observed a significant reduction in empty carcass weight of growing rabbit on diet with exogenous enzyme inclusion when maize replaced maize milling wastes. The results in this study showed that enzyme supplementation on the mash diet gave lower values of organ weights (as a percentage of the live weight) while the mash diet without supplementation gave the highest values suggesting that enzyme supplementation had a negative influence on organ weights. An explanation to this observation is that perhaps enzyme supplementation increased transit time, which made less nutrients available in the short run. Pelleting without enzyme inclusion led to increased ileum lengths, possibly a response to an increased surface area for the absorption of digesta. The reduced large intestine and colon lengths and the increased caecum of the rabbits on the sole could be a reflection of more undigested matter thus requiring more space for microbial degradation, to meet the animal's nutrient requirements. Al Bustany (1996) found that enzyme supplementation

was more effective on the mash than on the pelleted diets for improving the feed conversion ratio. Garcia *et al.* (2003) observed that using amylase enzyme of maize/soy diets improved feed efficiency, but it had no significant effect on carcass traits. Olayemi *et al.* (2006) however observed that carcass weight of growing rabbits was significantly reduced on diet with enzyme inclusion when replacing maize with maize milling waste. Iyayi and Davies (2005) however observed no significant difference in the organ weights of birds fed a PKC-enzyme supplemented diet and a control diet, though the kidney weight was observed to be significantly lower.

Mannanase inclusion on the mash diet had the greatest influence on GIT length. Diets with reduced fiber resulted in gut hypomotility, and prolonged retention time in the hindgut leading to an enlargement of the hindgut Cheeke (1994). Wang *et al.* (2005) had stated that the inclusion of exogenous enzymes significantly altered the morphology of the different GIT. NSPs bind to nutrients and reduce surface area for intestinal digestion. The presence of exogenous enzymes could hydrolyse complexes formed, thus reducing digesta viscosity and enabling better absorption of metabolites across the intestinal lumen. This could have been responsible for the better digestibility observed on diets containing mannanase. The influence of

supplementation of diets with exogenous enzymes varied according to different authors.

### Conclusion

This study demonstrated that enzyme supplementation and pelleting improved the carcass output. Pelleting positively influenced gastro-intestinal tract parameters but had a negative impact on mineral digestibility. Organs of rabbits fed enzyme supplemented diets showed reduction weights. Pelleting other than exogenous enzyme inclusion in the diets appeared effective means of improving rabbit performance however the cost of pelleting will add to the cost of production.

### References

- Adeniji A. A. 2008.** Replacement value of maize with enzyme supplemented decomposed bovine rumen content in the diets of weaner rabbits. *Asian Journal of Animal and Veterinary Advances*. 3 (2): 104-108.
- Al Bustany Z. 1996.** The effect of pelleting an enzyme supplemented barley-based broiler diet. *Anim. Feed Sci. Tech.*, 58: 283-288.
- Behnke, K. C. 1994.** Factors affecting pellet quality. Maryland Nutrition Conference. Dept. of Poultry Science and Animal Science, College of Agriculture, University of Maryland, College Park.
- Behnke K. C. 2001.** *Factors influencing pellet quality.* *FeedTech*. 5 (4) 19-22
- Cheeke, P. R. (1994).** Nutrition and nutritional diseases. In: P. J. Manning, D. H. Ringler and C. E. Newcomer. *The Biology of the Laboratory Rabbit*. 2<sup>nd</sup> ed. p 321. Academic Press, New York.
- Duncan, D. B. 1955.** Multiple range and multiple F-test. *Biometrics*, 11, 1-42.
- Falk D. (1985).** Feed Manufacturing Technology III. Ed. R.R. McElhiney. American Feed Industry Association Arlington, VA.
- FAOSTAT 2010.** Annual palm oil and palm kernel oil production in Nigeria. FAO estimates. FAO Statistics Division 2010. www.fao.org. Accessed 25 January 2010.
- Garcia M. I., M. J. R. Aran-bar., P. Medel and G. G. Mateous 2003.** Amylase supplementation of broiler diets based on corn. *Poultry Sci.*, 82: 436-442.
- Hathorn, D. J. and A. Mowlem 1980.** Some observations on pelleted diets for rabbits. *Journal of the Institute of Animal Technicians*, 31 (2): 69-76.
- Iyayi E. A. and B. I. Davies 2005.** Effect of enzyme supplementation of palm kernel meal and brewer's dried grain on the performance of broilers. *International Journal of Poultry Science*: 4(2): 76-80
- Longe O. G. and J. O. Fagbenro-Bryon (1990)** Composition and Physical Characteristics of some fibrous wastes and by-products of pig feeds in Nigeria. *Bieter. Trop. Lanwirtsch. Vet. Med.*, 28, 199-205.
- Han, Z. 1996.** Effect of enzyme supplementation of diets on the physiological function and performance of poultry. Proceedings of the 1<sup>st</sup> Chinese symposium on feed enzymes, Nanjing Agricultural University, Nanjing, China, 6-8 May 1996, pp 29-44
- Mosobalaje, M. A., O. A. Abu, Adedoyin, A. A. and O. O. Tewe 2012.** Bioconversion of palm kernel cake and its utilization in laying hens. Proceedings of 37<sup>th</sup> Annual of the Nigerian Society for Animal Production. Pp 250-252.
- Ng W. and K. Chong 2002.** The Nutritive Value of Palm Kernel Meal and the

Effect of Enzyme Supplementation in Practical Diets for Red Hybrid Tilapia (*Oreochromis sp.*) *Asian Fisheries Science* 15:167-176

**Olayemi W.A., A.O. Oso, A.M. Bamgbose, O.O. Oduguwa and S.A. Onadeko 2006.** Response of weaner rabbits to xylanase enzyme supplemented maize milling waste based diets. *Journal of Animal and Veterinary Advances* 5 (10): 839-843.

**Oluwafemi R. A. 2009.** Palm kernel Cake (PKC) utilization in Monogastric Animal Feeding Implications for Sustainable livestock Development. *The Internet Journal of Veterinary Medicine*. Volume 6 Number 2.

**Salari, S., H. Kermanshahi and H. Nasiri Moghadam 2006.** Isolation of a galactomannan that encourages macrophage activation from the edible *Morchella esculenta*. *J. Agric. Food Chem.* 50: 5683-5685

**Sarvestani T.S., N. Dabiri, M.J. Agah and H. Norollahi 2006.** Effect of Pellet and Mash Diets Associated with Biozyme Enzyme on Broilers Performance. *International Journal of Poultry Science* 5 (5): 485-490

**Smith W. T. 1997.** Commercial Rabbit Production. Mississippi State University Extension Service. Publication 1384. [www.msstate.edu](http://www.msstate.edu).

**SPSS 2001.** SPSS 11.0 for Windows. SPSS Inc. Chicago, USA.

**Sundu, B, A. Kumar and J. Dingle 2006.** Palm kernel meal in broiler diets: effect on chicken performance and health. *World Poultry Science Journal*, 62: 316-325.

**Wang Z. R., S. Y. Qiao, W. Q. Lu and D. F. Li 2005.** Effects of enzyme supplementation on Performance, nutrient digestibility, gastrointestinal morphology, and volatile fatty acid profiles in the hindgut of broilers fed wheat-based diets. *Poultry Science*, 84:875-881

**Yu B. and Tsen H.Y. 1993.** An *in vitro* assessment of several enzymes for the supplementation of rabbits' diets, *Animal Feed Science and Technology*; 40, 309-320.

**Ziggers D. 2004.** The better the pellet, the better the performance. *Feed Tech* 8 (6) 18-21.

Received: 15th April, 2012

Accepted: 10th August, 2012