

# *Gender differential in inclination to donate brain for research among Nigerians: the IBADAN Brain Bank Project*

**Rufus Akinyemi, Akin Ojagbemi, Joshua Akinyemi, Ayodeji Salami, Funmi Olopade, Temitope Farombi, Michael Nweke, Ezinne Uvere, et al.**

## **Cell and Tissue Banking**

International Journal for Banking, Engineering and Transplantation of Cells and Tissues Incorporating Advances in Tissue Banking


ISSN 1389-9333

Volume 20

Number 2

Cell Tissue Bank (2019) 20:297-306

DOI 10.1007/s10561-019-09769-4





# Gender differential in inclination to donate brain for research among Nigerians: the IBADAN Brain Bank Project

Rufus Akinyemi · Akin Ojagbemi · Joshua Akinyemi · Ayodeji Salami · Funmi Olopade · Temitope Farombi · Michael Nweke · Ezinne Uvere · Mayowa Aridegbe · James Balogun · Godwin Ogbole · Ayodele Jegede · Rajesh Kalaria · Adesola Ogunniyi · Mayowa Owolabi · Oyedunni Arulogun

Received: 14 December 2018 / Accepted: 17 April 2019 / Published online: 26 April 2019  
© Springer Nature B.V. 2019

## Abstract

**Background** Laboratory-based studies of neurological disease patterns and mechanisms are sparse in sub-Saharan Africa. However, availability of human brain tissue resource depends on willingness towards brain donation. This study evaluated the level of willingness among outpatient clinic attendees in a Nigerian teaching hospital.

**Methods** Under the auspices of the IBADAN Brain Bank Project, a 43—item semi-structured interviewer—

administered questionnaire was designed to evaluate knowledge, attitude, and beliefs of individuals attending Neurology, Psychiatry and Geriatrics Outpatient clinics regarding willingness to donate brain for research. Association between participants characteristics and willingness towards brain donation was investigated using logistic regression models. Analysis was conducted using Stata SE version 12.0.

**Results** A total of 412 participants were interviewed. Their mean age was 46.3 (16.1) years. 229 (55.6%)

R. Akinyemi · M. Aridegbe  
Institute for Advanced Medical Research and Training,  
College of Medicine, University of Ibadan, Ibadan,  
Nigeria

A. Ojagbemi  
Department of Psychiatry, College of Medicine,  
University of Ibadan, Ibadan, Nigeria

J. Akinyemi  
Department of Epidemiology and Medical Statistics,  
College of Medicine, University of Ibadan, Ibadan,  
Nigeria

A. Salami · M. Nweke  
Department of Pathology, College of Medicine,  
University of Ibadan, Ibadan, Nigeria

F. Olopade  
Department of Anatomy, College of Medicine, University  
of Ibadan, Ibadan, Nigeria

T. Farombi  
Chief Tony Anenih Geriatric Centre, University College  
Hospital, Ibadan, Nigeria

A. Jegede  
Department of Sociology, Faculty of the Social Sciences,  
University of Ibadan, Ibadan, Nigeria

R. Kalaria  
Institute of Neuroscience, Newcastle University,  
Newcastle upon Tyne, UK

E. Uvere · A. Ogunniyi · M. Owolabi  
Department of Medicine, College of Medicine, University  
of Ibadan, Ibadan, Nigeria

J. Balogun  
Department of Surgery, College of Medicine, University  
of Ibadan, Ibadan, Nigeria

G. Ogbole  
Department of Radiology, College of Medicine,  
University of Ibadan, Ibadan, Nigeria

were females and 92.5% had at least 6 years of formal education. Overall, 109 (26.7%) were willing to donate brains for research. In analyses adjusting for educational status, religion, ethnicity, marital status and family setting, male sex showed independent association with willingness towards brain donation OR (95% CI) 1.7 (1.08–2.69),  $p = 0.023$ . Participants suggested public engagement and education through mass media (including social media) and involvement of religious and community leaders as important interventions to improve awareness and willingness towards brain donation.

**Conclusion** The survey revealed low willingness among outpatient clinic attendees to donate brain for research, although men were more inclined to donate. It is imperative to institute public engagement and educational interventions in order to improve consent for brain donation for research.

**Keywords** Willingness · Brain donation · Brain banking · Nigeria · Africa · LMIC

## Introduction

African populations are ageing rapidly in keeping with trends in other low and middle income countries (LMIC). Concomitant with this is a projected increase in the prevalence of ageing—associated cerebrovascular and neurodegenerative disorders. Among the latter, Alzheimer's and Parkinson's diseases are relatively common (Akinyemi et al. 2014; Dotchin et al. 2013; Kalaria et al. 2008).

Post-mortem studies of the human brain are vital components of longitudinal research on ageing and neurodegenerative diseases for providing data to further understanding of prevalence, pattern, and mechanisms of disease (Allan et al. 2011; Bennett et al. 2006a, b). They are indeed critical to unravelling

associations with risk factors, discovering new treatment and preventative approaches, and exploring unique ethnic/racial neuropathological variations in brain structure and function (Barnes et al. 2015). However, laboratory-based histopathological and neurochemical studies of brain disorders are sorely lacking in sub-Saharan Africa due to lack of infrastructure, relevant skills and brain tissue resource from declining autopsy rates (Akinyemi et al. 2018a, b).

Biobanking is an emerging research field which is critical to personalized medicine (Hewitt 2011). With the declining autopsy rates in hospitals, organized brain donation initiatives have come up as an innovative intervention to ensure supply of brain tissue for relevant research, and they are often components of longitudinal studies (Kretzschma, 2009). Success, however, depends on optimal awareness of brain donation, autopsy, diagnostic and research utilization of brain tissue as well as willingness to provide consent among research participants, patients attending clinics for brain disorders and their caregivers, as well as the general public (Boise et al. 2017b; Jefferson et al. 2011, 2013).

In the last three decades, researchers in western countries have explored awareness and willingness towards brain tissue donation among volunteers in longitudinal ageing and dementia research programmes (Boise et al. 2017a, b; Jefferson et al. 2011, 2013). Findings from these studies suggest that educational attainment and culturally-shaped perceptions, attitudes, values and beliefs may influence willingness to participate in longitudinal brain research programmes or be a brain donor. Ethnic/racial backgrounds also often influenced willingness to participate in research and/or enlist as a potential brain donor (Boise et al. 2017a, b; Lambe et al. 2011).

Biobanking is growing in Africa but organized brain banking is virtually non-existent (Akinyemi et al. 2018a). Public understanding of brain biobanking in the context of unique culture, language, and belief systems has never been examined in Africa. The aim of this study was to determine willingness towards brain donation for research purpose among attendees of neuroclinics in an African teaching hospital. We hypothesized that there would be low willingness towards brain donation for research purpose among the study participants.

---

O. Arulogun  
Department of Health Promotion and Education, College of Medicine, University of Ibadan, Ibadan, Nigeria

R. Akinyemi (✉)  
Neuroscience and Ageing Research Unit, College of Medicine, University of Ibadan, Ibadan, Nigeria  
e-mail: rufusakinyemi@yahoo.com;  
roakinyemi@com.ui.edu.ng

## Methodology

### Setting

The IBADAN Dementia, Ageing and Neurodegeneration [IBADAN] Brain Bank Project is a multidisciplinary pilot project that aims to determine awareness and willingness to donate brain for research purpose, establish a pilot brain bank and evaluate the immunohistochemical features of ageing—associated degenerative and vascular pathologies in brain tissue from a pilot cohort of adult/older Nigerians (Akinyemi et al. 2018b). It is funded by the Research and Innovation Management Unit of the College of Medicine, University of Ibadan, Nigeria (Imam and Akinyemi 2016).

The current study was undertaken in the Neurology and Psychiatry Clinics as well as the Chief Tony Anenih Geriatric Centre (CTAGC) of the University College Hospital Ibadan (UCH), southwestern Nigeria. UCH is the premier academic teaching hospital in Nigeria that was established 60 years ago. It receives referrals from all over Nigeria and is well established as a center of excellence in Neurosciences, particularly stroke, ageing and dementia research. The CTAGC is the premier purpose built geriatric centre in Nigeria which caters for individuals 60 years and older.

### Sample

Using a cross-sectional study design, four hundred and twelve consenting adults aged 18–85 years were recruited from the outpatient clinics of the neurology, psychiatry and geriatrics departments of the University College Hospital Ibadan, Nigeria. Trained interviewers conducted face-to-face interviews with consenting study participants using a semi-structured questionnaire. Patients were identified during their routine presentation for follow-up at the respective Geriatrics, Neurology and Psychiatry clinics at the study site. Consenting patients who were adults (> 18 years of age); non-demented, mentally stable and capable of participating in discussion were recruited.

### Survey development

Survey data were collected using a 43-item interviewer administered semi-structured questionnaire which was categorized into sections including socio-demographic characteristics, awareness of brain disorders, willingness towards brain donation, and facilitators and barriers of brain donation. Questions on sociodemographic characteristics gathered include age, gender, race/ethnicity (Yoruba, Igbo, Hausa or other), marital status (single, married, widowed, separated, divorced), family setting (“monogamous”, “polygamous”) and educational status (no formal education, primary school, some secondary but failed to graduate, secondary school graduate, some university or 2-year degree, 4-year university graduate or more).

Awareness of brain diseases focused on perception of what brain diseases are and types of brain diseases. The statement ‘do you think brain disease can be cured?’ had responses coded as ‘Yes’ or ‘No’. This statement was followed by a probing question requesting open ended responses on ways in which brain diseases can be cured. These responses were later coded.

Respondents’ willingness towards brain donation for research purpose was assessed with a “yes” or “no” response. To highlight reasons for unwillingness towards brain donation, participants responded to the following statements: ‘I cannot donate my brain because of medical reasons’, “this is against my religion”, “I am afraid to donate because it is possible for a brain-dead person to recover from his/her injuries”, “I do not want my body cut up or disfigured”, “I don’t trust the health system”, “I don’t like the thought of it as such thoughts increase the chances of death”, “never thought about it”, “I want my body maintained whole as my parents gave birth to me”. Participants also responded to questions on why they were willing to consent to brain donation: “I think that after I die, I do not need my organs anymore”, “I like the idea that after I die, the findings from the examination of my brain can save someone’s life”, “I think if I die, my family would find solace in the idea that results of examination on my brain examination is helping someone else live”, and any other reasons.

The questionnaire was developed, revised, reviewed and presented to study investigators for their

comments and grammatical edits. This was subjected to another level of validation by experts in neurology, neurosurgery, pathology, mixed methods research who advised the team on arrangement, flow and presentation of questions to enhance eliciting responses from various cohorts.

After the survey instrument was finalized in English, it was translated into Yoruba language by a bi-lingual professional translator who worked in a community-based research support organization. Back translation was provided by a second bi-lingual professional translator at the same translation service. Discrepancies between the English version and the back translated version were reviewed and reconciled by an independent reviewer. The draft questionnaire was pretested among ten research subjects from a different patient population within the University College Hospital, Ibadan. Findings and comments from the pre-test were used to refine the questionnaire to ensure language appropriateness, coherence, clarity and reduction of completion time.

#### Ethical consideration

All procedures for data collection and analysis for the questionnaire survey phase of the project were reviewed and approved by the University of Ibadan/ University College Hospital Institutional Review Committee in Ibadan, Nigeria and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in the study.

#### Data collection procedures

Prior to commencement of data collection, a team comprising of eight field staff and volunteers were trained by experts from the Faculty of Public Health of the University of Ibadan on the requisite skills for collecting data using the semi-structured questionnaire and informed consent process.

Data collection commenced in the second week of June 2017 and lasted for a period of 3 weeks. The information was collected by the trained interviewers from qualified and consenting participant by 8 am on Mondays, Tuesdays, Thursdays and Fridays in a week at the Psychiatry, Neurology and Geriatric clinics respectively.

#### Data analysis procedure

Data from the completed copies of the questionnaire were entered and analyzed using the Stata SE version 12.0. Categorical variables were summarized using frequencies and percentages while mean (SD) was computed for continuous variables. Association between categorical variables and willingness towards brain donation was investigated using the Chi square test. Logistic regression models were fitted to estimated unadjusted and adjusted Odds Ratio (with their 95% Confidence Interval) for factors associated with willingness towards brain donation and adjusting for educational status, religion, ethnicity, marital status and family setting.

#### Results

A total of 412 participants completed the questionnaire survey of which 183 were males (44.4%) and 229 females (55.6%). Their mean age was 46.3 9 (SD = 16.1) years (Table 1). Thirty-one (7.5%) had no formal education while 50.2% attained tertiary education. The commonest religious affiliation and ethnic group were Christianity (70.6%) and Yoruba (92.7%) respectively. Majority were married (64.6%) and monogamy (79.1%) was the predominant form of marriage.

Three hundred and fifty-nine participants (87.1%) were aware of brain disease (Table 2). Among these, response to question on their perception about brain disease included: madness (13.3%), mental disorder (24.8%), brain malfunction (21.1%) and abnormality (21.1%). When asked about how they would approach the management of brain disease, medical treatment (66.0%) was the commonest response. Others were: spiritual intervention (20.1%) traditional healers (6.2%), psychiatrist attention (2.4%) and psychologist attention (4.1%).

Overall, 109 (26.7%) of respondents were willing to donate brains for research purpose. Table 3 shows the association between socio-demographic characteristics and willingness towards brain donation. There was no difference in the mean age of those willing (mean = 46.1 years) and not willing (46.4 years). Males (33.5%) were significantly more willing to donate brains than females (21.2%). Similarly, the proportion of those willing to donate increased

**Table 1** Sociodemographic characteristics of study participants (N = 412)

Variable	Total: 412 (100%)	Male: 183 (44.4%)	Female: 229 (55.6%)	<i>p</i> value
Age				
Mean age (years): mean (sd)	46.3 (16.1)	44.3 (16.8)	48.0 (15.4)	<b>0.0195*</b>
Education				
No formal education	31 (7.5)	5 (2.7)	26 (11.4)	
Primary	71 (17.2)	29 (15.9)	42 (18.3)	
Secondary	103 (25.0)	47 (25.7)	56 (24.5)	< <b>0.001*</b>
Some tertiary	85 (20.6)	31 (16.9)	54 (23.6)	
Graduate or more	122 (29.6)	71 (38.8)	51 (22.3)	
Religion				
Christianity	291 (70.6)	122 (66.7)	169 (73.8)	
Islam and others	121 (29.4)	61 (33.3)	60 (26.2)	0.114
Ethnic group				
Yoruba	382 (92.7)	169 (92.4)	213 (93.0)	
Non-Yoruba	30 (7.3)	14 (7.6)	16 (7.0)	0.066
Marital status				
Single	107 (26.0)	64 (35.0)	43 (18.8)	
Married	266 (64.6)	114 (62.3)	152 (66.4)	< <b>0.001*</b>
Widowed/separated/divorced	39 (9.4)	5 (2.7)	34 (14.9)	
Family setting				
Monogamous	326 (79.1)	157 (85.8)	169 (73.8)	
Polygamous	86 (20.9)	26 (14.2)	60 (26.2)	0.858

Asterisk and bold indicates statistically significant at  $p < 0.05$

progressively with higher educational attainment, although this failed to attain statistical significance. Willingness to donate brains did not vary across religious grouping, ethnicity and marital status. In contrast, participants from monogamous families were 1.9 times (OR 1.91, CI 1.04–3.49) more willing to donate brains than participants from polygamous settings. Awareness of brain disease was not associated with willingness towards brain donation. Multi-variable logistic regression revealed that male sex (OR 1.70, CI 1.08–2.69) was the main independent predictor of willingness to donate brains.

Table 4 summarizes the reasons for willingness (or lack of) towards brain donation for research purpose. The commonest reason for lack of willingness was that participants did not want their bodies disfigured after death (43.5%) while 39.4% never thought about it. The most common reason for willingness to donate brains was a respondent's knowledge that research done with his/her brain after death could help save lives (73.4%) (Table 5).

Participants were further asked about suggestions on how to promote brain donation. Suggestions provided were: mass media publicity (302, 73.8%), educational programmes (258, 63.1%), involvement of religious and community leaders (196, 47.5%), social media (151, 36.9%) and legislation (47, 11.5%).

## Discussion

This study which was aimed at investigating the barriers and facilitators of willingness to donate brain among Africans found that well over a quarter of participants were willing to donate brains with men having two-times the odds of willingness to donate brains. Participants in this study suggested public engagement using mass media and inclusion of religious and community leaders in these programmes as important interventions to improve inclination towards brain donation. Respondents who were willing to donate brain for research gave such reasons as

**Table 2** Awareness, perception and knowledge of brain disease

Item	Total: 412 (100%)	Male: 183 (44.4%)	Female: 229 (55.6%)	<i>p</i> value
Aware of brain disease	359 (87.1)	161 (88.0)	198 (86.5)	0.648
Perception about about brain disease <sup>a</sup>				
Madness	53(13.3)	21 (13.5)	32 (17.3)	0.452
Mental disorder	99 (24.8)	46 (29.7)	53 (28.6)	0.638
Brain malfunction	84 (21.1)	45 (29.0)	39 (21.1)	0.058
Neurological disorder	6 (1.5)	5 (3.2)	1 (0.5)	0.093f
Loss of memory	8 (2.0)	8 (5.2)	0 (0.0)	<b>0.001*f</b>
Mental retardness	7 (1.8)	3 (1.9)	4 (2.2)	1.000
Brain damage	10 (2.25)	4 (2.6)	6 (3.2)	1.000
Abnormality	84 (21.1)	30 (19.4)	54 (29.2)	0.072
Disease in the brain	22 (5.5)	10 (6.5)	12 (6.5)	0.919
Others	26 (6.5)	13 (8.4)	13 (7.0)	0.554
Approach to management of brain diseases				
Medical treatment	309 (66.0)	134 (63.8)	175 (67.8)	0.457
Spiritual intervention	94 (20.1)	44 (21.0)	50 (19.4)	0.595
Traditional healers	29 (6.2)	9 (4.3)	20 (7.8)	0.104
Psychiatrist attention	11 (2.4)	9 (4.3)	2 (0.8)	<b>0.027*f</b>
Psychologist attention	5 (1.1)	3 (1.4)	2 (0.8)	0.663f
Care from relations	19 (4.1)	11 (5.2)	8 (3.1)	0.346f

f: Fisher Exact test

<sup>a</sup>Multiple response itemsAsterisk and bold indicates statistically significant at  $p < 0.05$ 

the fact that they don't need the brain or other organs after death, research done using their brains might contribute to medical progress and saving others and that their families would love the idea that they donated brain for research. Barriers to brain donation identified by the respondents include fear of body disfigurement from autopsy, mistrust of the health system, religious belief, and belief that a brain-dead person might recover.

A high level of awareness about brain diseases was expected in this study sample given the fact that a very good proportion was recruited from neuroclinics. The varying perceptions expressed regarding brain diseases however could be due to the differing levels of education, and the religious and cultural perspectives held by respondents regarding the sort of conditions on account of which the respective clinics were visited. Indeed, cultural beliefs and practices particularly influence the approach to the diagnosis and management of brain disorders in sub-Saharan African climes (Adekson 2016; Osuntokun 1975). These factors also influenced the suggested approaches to the treatment

of brain disorders wherein spiritual treatment was suggested following medical treatment. Respondents also suggested the involvement of community and religious leaders as measures to improve awareness towards brain donation. The strong influence of religious beliefs on the perceived causation of illnesses, health choices and practices have been variously explored in diverse settings among indigenous Africans (Hurst et al. 2015; Jenkins et al. 2016), African Americans (Jefferson et al. 2011; Lambe et al. 2011) and Caucasians (Jefferson et al. 2011).

Only about one third of respondents in this study indicated willingness to donate brains for research purposes despite the claims of high level of awareness of brain disorders. Willingness to donate was recorded in less than 50% of elderly African Americans whereas 75% of elderly Caucasians consented in a previous study (Jefferson et al. 2011). Although we have specifically explored ethnicity in this current study, evidence from literature suggest that minority populations of African ancestry tend to exhibit lower willingness towards brain donation compared to their

**Table 3** Association between socio-demographic characteristics and willingness towards brain donation

Variable	Yes	No	Unadjusted			Adjusted		
			OR	95% CI	<i>p</i> value	OR	95% CI	<i>p</i> value
Age (years): mean (SD)	46.1 (16.5)	46.4 (16.1)	0.99	0.98–1.01	0.895			
Sex								
Male	61 (33.5)	121 (66.5)	1.88	1.21–2.93*	<b>0.005*</b>	1.7	1.08–2.69	<b>0.023*</b>
Female	48 (21.2)	179 (78.8)	1.00			1.00		
Education								
No formal education	4 (13.3)	26 (86.7)	1.00			1.00		
Primary	17 (23.9)	54 (76.1)	2.05	0.63–6.69	0.236	1.65	0.49–5.49	0.415
Secondary	28 (27.5)	74 (72.6)	2.46	0.79–7.68	0.121	1.84	0.58–5.90	0.303
Some tertiary	23 (27.1)	62 (72.9)	2.41	0.76–7.66	0.136	1.88	0.58–6.09	0.294
Graduate or more	37 (30.6)	84 (69.4)	2.86	0.93–8.79	0.066	1.89	0.59–6.06	0.282
Religion								
Christianity	79 (27.4)	209 (72.6)	1.15	0.70–1.87	0.582			
Islam and others	30 (24.8)	91 (75.2)	1.00					
Ethnic group								
Yoruba	102 (26.9)	277 (73.1)	1.21	0.51–2.91	0.67			
Non-Yoruba	7 (23.3)	23 (76.7)	1.00					
Marital status								
Single	32 (29.9)	75 (70.1)	1.00					
Married	68 (25.8)	196 (74.2)	0.81	0.49–1.34	0.415			
Widowed/separated/divorced	9 (23.7)	29 (76.3)	0.73	0.31–1.71	0.465			
Family setting								
Monogamous	94 (29.0)	230 (71.0)	1.91	1.04–3.49	<b>0.037*</b>	1.62	0.86–3.04	0.137
Polygamous	15 (17.7)	70 (82.4)	1.00			1.00		
Awareness of brain disease	99 (27.7)	258 (72.3)	1.61	0.78–3.34	0.199			

Asterisk and bold indicates statistically significant at  $p < 0.05$

Caucasian counterpart (Boise et al. 2017b; Jefferson et al. 2013; Lambe et al. 2011).

Understanding that the brain and other body organs are no more needed after death and appreciation of the need for contribution to research in order to facilitate new discoveries and medical progress have been identified as key facilitators of willingness to donate brain for research in previous studies (Jefferson et al. 2013; Lambe et al. 2011; Oluwasola et al. 2009). This is usually a key message of brain donation enlistment programmes when volunteers join longitudinal research projects or during public engagement programmes related to such researches (Arulogun et al. 2016).

Fear of disfigurement has been identified in several studies as a barrier to donation of brain and other organs (Lambe et al. 2011; Oluwasola et al. 2009). It

has its foundations in cultural and religious beliefs and practices that a deceased's body needs to be kept 'whole' in order to enable him function well in the after life, upon re-incarnation or because of religious demands (Boise et al. 2017a, b). A similar reason was found as a barrier to autopsy among Nigerian patients and caregivers in a previous study (Oluwasola et al. 2009). On the other hand, mistrust of the health system could be a reaction to perceived wrongs suffered in the past (Lambe et al. 2011) or an expression of dissatisfaction of quality of healthcare received by the deceased person while alive.

Educational programmes using the traditional media have been used with success to enlighten the public and patient populations and raise awareness about health issues including the benefit of participating in research involving donation of biological

**Table 4** Reasons for willingness (or lack of) towards brain donation for research purpose

Reasons	Total: n (%)	Male: n (%)	Female: n (%)	<i>p</i> value
Lack of willingness to donate brain (n = 300) <sup>a</sup>				
Medical reasons	13 (5.1)	7 (7.5)	6 (3.8)	0.312
Religion	35 (13.8)	11 (11.7)	24 (15.1)	0.254
Afraid because brain-dead person can recover	17 (6.7)	9 (9.6)	8 (5.0)	0.276
Don't want body disfigured	110 (43.5)	40 (42.6)	70 (44.0)	0.288
Don't trust medical system	24 (9.5)	11 (11.7)	13 (8.2)	0.567
Thoughts about higher chances of death	34 (13.4)	14 (14.9)	20 (12.6)	0.915
Never thought about it	100 (39.4)	38 (40.4)	62 (38.8)	0.56
I want my body whole	34 (13.9)	12 (13.2)	22 (14.4)	0.525
Reason for willingness to donate brain (n = 109) <sup>a</sup>				
I don't need it after death	60 (61.9)	36 (65.5)	24 (57.1)	0.348
Findings from my brain examination can save a life	80 (82.5)	43 (78.2)	37 (88.1)	0.439
My family will be pleased that examination of my brain safe life	33 (34.7)	25 (46.3)	8 (19.5)	<b>0.006*</b>

<sup>a</sup>Multiple response items

Asterisk and bold indicates statistically significant at  $p < 0.05$

**Table 5** Suggestions to promote willingness to donate brain for research purpose (n = 409)

Suggestion	Total: n (%)	Male: n (%)	Female: n (%)	<i>p</i> value
Media publicity	302 (73.8)	144 (79.1)	158 (69.6)	0.131
Education	258 (63.1)	128 (70.3)	130 (57.3)	<b>0.028*</b>
Legislation	47 (11.5)	24 (13.2)	23 (10.1)	0.417
Involve religious and community leaders	196 (47.9)	92 (50.6)	104 (45.8)	0.578
Educate people on social media	151 (36.9)	83 (45.6)	68 (30.0)	<b>0.003*</b>

Multiple responses

Asterisk and bold indicates statistically significant at  $p < 0.05$

materials in life or death (Akinyemi et al. 2018a; Jefferson et al. 2013; Jenkins et al. 2016; Moodley and Singh 2016). Engaging community and religious leaders has also been an effective strategy in enhancing research participation and dealing with issues unique to specific populations with peculiar beliefs and practices (Wells et al. 2004; Wight et al. 2016). This strategy, in particular, has been useful in engaging minority population such as African Americans in the United States (Jefferson et al. 2013). Social media, mobile telephony and other emerging mHealth platforms (Sarfo et al. 2016) are particularly useful approaches for reaching the younger generation, and will be strategic for the future of neurobiobanking initiatives (Akinyemi et al. 2018b).

#### Strengths, limitations and future directions

This study, to the best of our knowledge, represents the first attempt to study willingness towards brain donation for neurobiobanking in Nigeria with implications for other low and medium income countries (LMICs) (Akinyemi et al. 2018b). The study process provided an opportunity to engage potential brain donors for the Ibadan Brain Bank Project. Individuals attending neuroclinics were studied, hence the findings may not be generalizable to the general population with non-neurological conditions or disorders. This could have potential implications for the accrual of control brains in a neurobiobank, but hopefully public engagement programmes suggested by the study participants could also enhance awareness and inclination towards brain donations in the general

population without neurological morbidities. Nevertheless, findings from this study offer unique insight into the barriers and facilitators of willingness towards brain donation and provides strategies for developing intervention strategies to educate and engage potential brain donors to enlist in programmes for establishing neurobiobanks in sub-Saharan Africa.

**Acknowledgements** The IBADAN Brain Bank Project is supported by Grant CTR16A012 from the College of Medicine, University of Ibadan, Nigeria while the African Neurobiobank ELSI Project, the SIREN Study and the Systematic Investigation of Blacks with Stroke using Genomics (SIBS Genomics) Study are supported by Grants U01HG010273, U54HG007479 and R01NS107900 respectively from the National Institutes of Health (NIH) as part of the H3Africa Consortium. RA is further supported by an IBRO Return Home Fellowship Grant. RNK acknowledges the support of the Medical Research Council, UK (G1100540) to the Newcastle Brain Tissue Resource (NBTR).

**Funding** This study was funded by Grant CTR16A012 from the College of Medicine, University of Ibadan, Ibadan, Nigeria.

#### Compliance with ethical standards

**Conflict of interest** All the authors declare no conflict of interest.

**Ethical approval** All procedures performed involving human participants in this study were in accordance with the ethical standards of the University of Ibadan/University College Hospital Institutional Review Committee in Ibadan, Nigeria and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

**Informed consent** Informed consent was obtained from all individual participants included in the study.

#### References

- Adekson MO (2016) Similarities and Differences Between Yoruba Traditional Healers (YTH) and Native American and Canadian Healers (NACH). *J Relig Health* 55:1717–1728
- Akinyemi RO et al (2014) Contribution of noncommunicable diseases to medical admissions of elderly adults in Africa: a prospective, cross-sectional study in Nigeria, Sudan, and Tanzania. *J Am Geriatr Soc* 62:1460–1466
- Akinyemi RO et al (2018a) Biobanking in a challenging African environment: unique experience from the SIREN Project. *Biopreserv Biobank*. <https://doi.org/10.1089/bio.2017.0113>
- Akinyemi RO et al (2018b) Brain banking in low—and middle-income countries: Raison D'être for the Ibadan Brain Ageing, Dementia and Neurodegeneration (IBADAN) Brain Bank Project. *Brain Res Bull* 145:136–141
- Allan LM et al (2011) Long term incidence of dementia, predictors of mortality and pathological diagnosis in older stroke survivors. *Brain* 134:3716–3727
- Arulogun OS et al (2016) Experience of using an interdisciplinary task force to develop a culturally sensitive multi-pronged tool to improve stroke outcomes in Nigeria. *eNeurologicalSci* 4:10–14
- Barnes LL et al (2015) Mixed pathology is more likely in black than white decedents with Alzheimer dementia. *Neurology* 85:528–534
- Bennett DA et al (2006a) Neuropathology of older persons without cognitive impairment from two community-based studies. *Neurology* 66:1837–1844
- Bennett DA, Schneider JA, Tang Y, Arnold SE, Wilson RS (2006b) The effect of social networks on the relation between Alzheimer's disease pathology and level of cognitive function in old people: a longitudinal cohort study. *Lancet Neurol* 5:406–412
- Boise L, Hinton L, Rosen HJ, Ruhl M (2017a) Will my soul go to heaven if they take my brain? Beliefs and worries about brain donation among four ethnic groups. *Gerontologist* 57:719–734
- Boise L et al (2017b) Willingness to be a brain donor: a survey of research volunteers from 4 racial/ethnic groups. *Alzheimer Dis Assoc Disord* 31:135–140
- Dotchin CL, Akinyemi RO, Gray WK, Walker RW (2013) Geriatric medicine: services and training in Africa. *Age Ageing* 42:124–128
- Hewitt RE (2011) Biobanking: the foundation of personalized medicine. *Curr Opin Oncol* 23:112–119
- Hurst S et al (2015) Pretesting qualitative data collection procedures to facilitate methodological adherence and team building in Nigeria. *Int J Qual Methods* 14:53–64
- Imam I, Akinyemi R (2016) Nigeria. *Pract Neurol* 16:75–77
- Jefferson AL, Lambe S, Cook E, Pimontel M, Palmisano J, Chaisson C (2011) Factors associated with African American and White elders' participation in a brain donation program. *Alzheimer Dis Assoc Disord* 25:11–16
- Jefferson AL, Lambe S, Romano RR, Liu D, Islam F, Kowall N (2013) An intervention to enhance Alzheimer's disease clinical research participation among older African Americans. *J Alzheimers Dis* 36:597–606
- Jenkins C et al (2016) Stroke investigative research and education network: community engagement and outreach within phenomics core. *Health Educ Behav* 43:82S–92S
- Kalaria RN et al (2008) Alzheimer's disease and vascular dementia in developing countries: prevalence, management, and risk factors. *Lancet Neurol* 7:812–826
- Kretzschmar H (2009) Brain banking: opportunities, challenges and meaning for the future. *Nat Rev Neurosci* 10:70–78
- Lambe S, Cantwell N, Islam F, Horvath K, Jefferson AL (2011) Perceptions, knowledge, incentives, and barriers of brain donation among African American elders enrolled in an Alzheimer's research program. *Gerontologist* 51:28–38
- Moodley K, Singh S (2016) "It's all about trust": reflections of researchers on the complexity and controversy surrounding biobanking in South Africa. *BMC Med Ethics* 17:57
- Oluwasola OA, Fawole OI, Otegbayo AJ, Ogun GO, Adebamowo CA, Bamigboye AE (2009) The autopsy:

- knowledge, attitude, and perceptions of doctors and relatives of the deceased. *Arch Pathol Lab Med* 133:78–82
- Osuntokun BO (1975) The traditional basis of neuropsychiatric practice among the Yorubas of Nigeria. *Trop Geogr Med* 27:422–430
- Sarfo FS et al (2016) Phone-based Intervention under Nurse Guidance after Stroke (PINGS): study protocol for a randomized controlled trial. *Trials* 17:436
- Wells K, Miranda J, Bruce ML, Alegria M, Wallerstein N (2004) Bridging community intervention and mental health services research. *Am J Psychiatry* 161:955–963
- Wight D, Wimbush E, Jepson R, Doi L (2016) Six steps in quality intervention development (6SQuID). *J Epidemiol Community Health* 70:520–525

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

UNIVERSITY OF IBADAN LIBRARY