

Microbiological Assessment and Detection of Adenovirus in Sachet Water Sold In Abeokuta, Nigeria

Shittu¹, O.B., Adekunle, O.T.,¹ Olufemi, F.O.,² Faleye, T.O.C.,³ Adewumi³, M.O. and Adeniji³, J.A.

1. Department of Microbiology, College of Biosciences, Federal University of Agriculture, Abeokuta, Nigeria.
2. Department of Veterinary Microbiology and Parasitology, College of Veterinary Medicine Federal University of Agriculture, Abeokuta, Nigeria.
3. Department of Virology, College of Medicine, University of Ibadan, Ibadan, Nigeria.

Abstract: Microbiological safety of sachet water remains a public health problem in Nigeria. This study was aimed at investigating some packaged sachet water sold in Abeokuta, South-West Nigeria for the microbiological safety including some of the enteric viruses on contaminant candidate list. Sachet water samples from five different producers were obtained over three month's period. Bacterial and fungal analyses were conducted with standard culture method. Targeted protozoans were investigated by microscopic examination of sediments obtained after centrifugation. Nested and semi-nested polymerase chain reaction (PCR) techniques targeting specific genes in adenovirus, norovirus and rotavirus were used for viral analyses. Results were presented in presence-absence score. Contingency table was used to establish relationship between viruses, *Escherichia coli* and protozoans. Out of a total twenty pooled samples analysed, adenovirus had a prevalence rate of 10% across the study period, whereas rotavirus and norovirus were absent. *Giardia* cysts and *Cryptosporidium* oocysts were also absent. *Escherichia coli* was present in 40% of the brands. Other bacteria identified were *Salmonella enterica* serovar Typhi, *Shigella dysenteriae*, and *Pseudomonas aeruginosa*. *Aspergillus* sp, *Mucor* and *Rhizopus* sp. were present in some samples collected. Adenovirus was detected by PCR in a pooled sample of sachet water that tested negative for *Escherichia coli*, *Cryptosporidium* oocysts and *Giardia* cysts. There is need for microbiological screening of sachet water periodically in order to enhance public health safety.

Key words: Adenovirus, Fungi, Norovirus, Protozoans, Rotavirus, Sachet water

INTRODUCTION

Poor water quality is associated with an estimated 3.5 billion diarrhoeal episodes and 1.87 million diarrhoeal-associated childhood deaths annually (Arnold and Colford, 2007; Boschi-Pinto *et al.*, 2008). Of these deaths, 90% occur in children from developing countries and this high proportion accounted for nearly 20% of the 10 million total deaths per annum in children under 5 years of age (Boschi-Pinto *et al.*, 2008; UNICEF, 2008). Pathogenic bacteria, viruses and protozoans are well known microbial contaminants of drinking water (Szewzyk *et al.*, 2000), although fungi are considered emerging chronic water quality problem (Hageskal *et al.* 2009; Ashbolt, 2015).

Waterborne viral infection is one of the most important causes of human morbidity (Fongaro *et al.*, 2013). Waterborne viruses have gained attention worldwide as

emerging pathogens because of their low infectious dose, survival in water and considerable health impacts (Swenson *et al.*, 2003; Fong and Lipp 2005; Xagorarakis *et al.*, 2007). As part of the Safe Drinking Water Act, United State Environmental Protection Agency (USEPA) listed adenoviruses and noroviruses as two of the four viral groups on the "Contaminant Candidate List" (CCL) (Miagostovich *et al.*, 2008; Teunis *et al.*, 2008; USEPA, 2009). In Nigeria, many households in rural and urban areas consume packaged sachet water (Odeyemi, 2015) because it is cheap, affordable, and readily available and also because of its perceived safety (Dada, 2009). There have been several studies on bacteriological quality of different brands of sachet water marketed in Nigeria using bacterial indicators without cognizance of virological quality.

Absence of bacterial indicators of fecal contamination does not necessarily guarantee consumers' safety from enteric pathogens like viruses (USEPA, 1998; Xagorarakis *et al.*, 2007; Hssaine *et al.*, 2011; Armon, 2015) and due to the cost of analysis, frequent virus monitoring programmes are currently impractical (Matthijnssens *et al.*, 2008). Viral contamination is best monitored by direct detection of the pathogens themselves without using indicators as a proxy (Armon, 2015). This study was therefore designed to assess some brands of sachet water sold in Abeokuta, South-West Nigeria for the presence of pathogens including some of the viruses on contaminant candidate list.

MATERIALS AND METHODS

Study Area

Abeokuta is one of the most prominent urban settlements in the SouthWestern Nigeria (Bello and Falano, 2017). It is the capital of Ogun State, lying between latitude 7° 06' and 7° 13' North and longitude 3° 15' and 3° 25' East (Olowofela *et al.*, 2013). Situated within the rainforest belt of the tropics, the city occupies a geographical area of 1256sqkm with a population of about 449,088 inhabitants (National Population Commission, 2006). The city is approximately 100km north of Lagos and 80km Southwest of Ibadan, the Oyo State capital. Inadequate public water supply is a major problem in the city (Odjegba *et al.*, 2015).

Sachet water sampling

A total of 60 sachet water samples were collected from five different producers. Water samples for viral analyses were frozen while others were stored at 4°C and analysed within 6 h of collection. Samples were pooled into a total of twenty (20) for analyses. Samples for viral analyses were transported to the virology laboratory of the Department of Virology, College of

Medicine, University of Ibadan, Ibadan, Nigeria.

Microbiological analyses

Water samples were inoculated by spread plate method on molten nutrient agar and incubated at 37°C for 24 h for total heterotrophs, MacConkey agar incubated at 37°C for 24 h for coliforms, and eosin methylene blue (EMB) agar incubated at 44°C for 24 h for *Escherichia coli*. Gram's staining and biochemical tests such as oxidase, catalase, sugar fermentation, urease reaction, lysine decarboxylase, indole test, and H₂S production were performed to identify bacterial isolates using Bergey's Manual of Systematic Bacteriology (Garrity *et al.*, 2004).

Fungi were detected by inoculating water samples on Sabouraud dextrose agar (SDA) and incubated at 25 °C for 7 days (Gottlich *et al.*, 2002). Microscopic identification was based on morphological characteristics of spores, macroconidia, microconidia, rhizoids and stolons according to Barnett and Hunter (2006).

Cryptosporidium parvum and *Gardia lamblia* were detected by repeated centrifuging of ten 50ml of each sample at 2,500 rpm for 5min. and the sediments obtained were subjected to microscopic examination using x 10 and x 40 objective lenses (Kwakye-Nuako *et al.*, 2007; Chinyelu *et al.*, 2010).

Viral analysis was carried out by following concentration protocol as described by WHO, (2003) and Hsainne *et al.* (2011). Sachet water (500ml of pooled samples) was collected and 39.5 ml of 22% dextran, 287ml 29% PEG6000, and 35 ml 5N NaCl were added. It was mixed thoroughly and kept in constant agitation for 1 h at 4°C using a horizontal shaker. Afterwards, the mixture was transferred into a sterile one litre separation funnel attached to retort stand and left overnight at 4°C. In the morning, 5.0 to 10 ml of lower layer was collected into sterile 50 mL centrifuge tube and stored at -20°C until analysed.

Viral nucleic acid was extracted using Qiagen viral mini kit (Qiagen, Germany) according to manufacturer's instructions. For rotavirus and norovirus cDNA synthesis, SCRIPT cDNA synthesis kit (Jena Bioscience, Germany) was used according to manufacturer's instruction. Briefly, for a 20.0µL cDNA mix, 12.0µL of extract was added to 0.5µL SCRIPT reverse-transcriptase, 4.0µL of SCRIPT RT buffer complete, 1.0µL dNTP mix, 1.0µL of DTT, 0.5µL of random hexamers, and 1.0µL RNase inhibitor. The reaction was incubated at 42°C for 10 minutes followed by 50°C for 60 minutes in a Veriti Thermal cycler (Applied Biosystems, USA). The cDNA was stored at -80°C until analysed.

PCR procedure for viral gene amplification

Target genes, and primers for viral gene amplification are presented in Table 1. PCR was done in 25µL volumes containing 5.0µL of Red load Taq (Jenabioscience), 5.0µL of Nucleic acid extract (for the adenovirus screen) or cDNA (for norovirus and rotavirus), 1µL of each primer and 13µL of RNase free water for the first round of PCR. Thermal cycling was 94°C for 3 min, 45 cycles at 94°C for 30 s, 56°C for 30 s and 72°C for 30 s. This was followed by 72°C for 7 min and held at 4°C till terminated. Amplicons from the first round of PCR were used as template for the second set of primers and the second PCR. All PCR assays were executed in a Veriti Thermal cycler (Applied Biosystems, USA) and all PCR products were resolved on 2% agarose gels stained with ethidium bromide and viewed using a transilluminator.

RESULTS

Detection of microorganisms of interest found in sachet water brands

Bacteria, fungi and viruses detected in sachet water samples are presented in Table 2. *Escherichia coli* was not detected in brands 1, 4 and 5 within the period of study while in brands 2 and 3, *Escherichia coli* Also, fungal analysis revealed varying levels of predominance of *Aspergillus sp.* (50%),

was detected. Fig. 1 shows *Escherichia coli* (17%), *Salmonella enterica* serovar Typhi (14%), *Pseudomonas aeruginosa* (3%), *Enterobacter aerogenes* (6%), *Shigella dysenteriae* (11%), *Staphylococcus aureus* (20%), and *Klebsiella aerogenes* (29%). Fig. 2 shows the occurrence of *Aspergillus niger* (50%), *Rhizopus oryzae* (40%) and *Mucor mucorales* (10%) in the sachet water sampled. Also, no ova, cyst, oocyst, trophozoite, or adult form of any protozoans was seen in all samples.

For enteric viruses, only 2 sample tested positive for adenovirus over the study period. Rotavirus and norovirus were not detected in all water samples collected. A cross tabulation of relationship between viruses (norovirus, adenovirus and rotavirus) examined versus the protozoans showed a close relationship of 100%, 93%, 100% respectively. The absence of these protozoans was observed to consistently indicate the absence of rotavirus, and norovirus in sachet water being investigated. *Escherichia coli* on the other hand indicated poor relationship to rotavirus, adenovirus and norovirus (60%, 57% and 60%) respectively (Table 3).

DISCUSSION

Water borne diseases remain a challenge in both developed and developing countries. Infectious diseases predominantly caused by human and animal enteric pathogens and health risks associated with drinking of non-potable water are well documented (WHO/UNICEF, 2006; Reynolds *et al.*, 2008; WHO, 2008). The bacteriological analysis in this study has revealed that some sachet water were heavily contaminated with persistent occurrence of *P. aeruginosa* and *S. dysenteriae*. The presence of *P. aeruginosa* and *Shigella dysenteriae* in some vended sachet water has been reported from various parts of the country (Adekunle *et al.*, 2004; Ezeugwunne *et al.*, 2009; Oladipo *et al.*, 2009; Shittu *et al.*, 2013, 2014; Mbah and Muhammed, 2015).

Rhizopus sp. (40%) and *Mucor sp.* (10%). Fungi have been reported from sachet and bottled drinking water (Shittu *et al.*, 2016; Thliza *et al.* 2015; Jonathan *et al.*, 2016). The presence of filamentous fungi in drinking water has become an area worthy of investigation especially with respect to biofilm formation and problems associated with pathogenic fungi and mycotoxins (Siqueira *et al.*, 2011; Paterson and Lima 2015; Oliveira *et al.* 2016; Novak Babič *et al.*, 2017; 2018; Paterson, 2019; Mhlongo *et al.*, 2019).

In this study, no oocyst of *Cryptosporidium parvum* or cyst of *Giardia lamblia* was detected in any sample including where adenovirus was found present. The absence of these protozoans could possibly be due to the low sensitivity of the microscopy method used. Molecular detection techniques such as PCR-based methods offer many advantages over microscopic method being rapid, specific, and sensitive (Yu *et al.*, 2009; Gotfred-Rasmussen *et al.*, 2016). Oocyst of *Cryptosporidium parvum* and cyst of *Giardia lamblia* have been reported in finished water (LeChevallier *et al.*, 1991; Kwakye-Nuako *et al.*, 2007; Chinelu *et al.*, 2010).

The detection of adenovirus in drinking water in this study was consistent with similar studies on tap water and river water (Cho *et al.*, 2000), South-African waters (Genthe *et al.*, 1995), raw and treated water (Van Heerden *et al.*, 2003; 2004); swimming pool (Van Heerden *et al.*, 2005), surface water and drinking water resources in Southern Ghana (Gibson-Schwab *et al.*, 2011) and in drinking water sources used in rural areas of Benin, West Africa (Verheyen *et al.* (2009). Human adenoviruses (HAdVs) are the second-leading cause of childhood gastroenteritis worldwide (USEPA, 2005). These are important human pathogens and are responsible for both enteric illnesses and respiratory and eye infections, acute hemorrhagic cystitis, and meningoencephalitis (Mena and Gerba, 2009).

Escherichia coli as a faecal indicator was not detected in the sample that was positive for adenovirus similarly to the study of Pusch *et al.* (2005) in German environmental waters. Adenoviruses have been found to be significantly more stable than faecal indicator bacteria and other enteric viruses during UV treatment (Jiang *et al.*, 2006). Enteric viruses have been isolated from water that indicated microbiological quality. Also, outbreaks of viral gastroenteritis have been reported from ingestion of water that complied with faecal coliform standard (Fong and Lipp, 2005).

From this study, absence of norovirus was in contrast to report of (Jack *et al.*, 2013). Noroviruses (NoVs), previously called Norwalk-like viruses, cause gastroenteritis in all age groups (Maunula *et al.*, 2005) and are the most common cause of acute nonbacterial gastroenteritis worldwide (Karim *et al.*, 2004; Blanton *et al.*, 2006). Noroviruses similar to other enteric viruses and can remain infectious in the environmental waters for long periods, surviving longer than bacteria (Green, 2007; Maunula, 2007; Teunis *et al.*, 2008; Seitz *et al.*, 2011).

Also, rotavirus was not detected in this study contrary to Verheyen *et al.* (2009) who found both rotaviruses and adenoviruses in drinking water. Rotavirus (RoV) is the most common cause of diarrhoeal disease primarily in young children less than five worldwide, though infection and disease in older children and adults also occur (Kapikian *et al.*, 2001; Bernstein 2009; Matthijnssens *et al.*, 2008). By the age of five, nearly every child in the world has been infected with rotavirus at least once (USEPA, 2005).

Though norovirus and rotavirus were not found in this study, both have been reported in clinical samples alongside with adenovirus from Southwest Nigeria (Arowolo *et al.*, 2019). This study has provided information on the occurrence of human enteric viruses in water from SW, Nigeria, relevant to research priorities on

human enteric viruses in Africa (Upfold *et al.*, 2021).

CONCLUSION

Public health significant pathogenic bacteria, opportunistic fungi and adenovirus were detected in packaged sachet water sold in

Abeokuta. This is indicative of potential microbiological hazards and risks especially, to children and immunocompromised individuals. Packaged sachet water in Nigeria requires increased attention and monitoring by the regulatory agencies.

Table 1. Target genes, PCR assay and primer sequences for viral analyses

Target Genes and PCR assay	Primers	Sequences	References
<i>Adenovirus hexon gene</i>	Forward 1	JTVXF (5'-GGACGCCTCGGAGTACCTGAG-3')	Xagorarakhi <i>et al.</i> (2007).
	Reverse 1	JTVXR (5'-ACIGTGGGGTTTCTGAACTTGTT-3')	
Semi-nested PCR	Forward 2	JTVXP (5'-CTGGTGCAGTTCGCCCCGTGCCA-3')	Kageyama <i>et al.</i> (2003)
	Forward 2	JTVXR (5'-ACIGTGGGGTTTCTGAACTTGTT-3')	
<i>Norovirus ORF gene</i>	Forward 1	JJGII (5'-CAAGAGTCAATGTTTAGGTGGATGAG-3')	Miagostovich <i>et al.</i> (2008)
	Reverse 1		
Nested PCR	Forward 2	GOG2R (5'-TCGACGCCATCTTCATTCACA-3')	
	Forward 2	RingP (5-TGGGAGGGCGATCGCAATCT-3) GOG2R (5-TCGACGCCATCTTCATTCACA-3)	
<i>Rotavirus VP7 gene</i>	Forward 1	ROTA VP7F1 (5'-GGCTTTAAAAGAGAGAATTTC-3')	
	Reverse 1		
Nested PCR	Forward 2	ROTA VP7R1 (5'-GGTCACATCATAACAATTCT-3')	
	Forward 2	ROTA VP7F2 (5'-TAGCTCCTTTTAATGTATGG-3') ROTA VP7R2 (5'-AACTTGCCACCATYTYTTCC-3')	

Table 2: Detection of microorganisms of interest found in sachet water brands

Microorganisms	Packaged sachet water brands				
	1	2	3	4	5
Bacteria					
<i>Escherichia coli</i>	---	+++	+++	---	---
Fungi					
<i>Aspergillus niger</i>	---+	---	++-	-+-	-+-
<i>Rhizopus oryzae</i>	---	---	+_-+	-++	---
<i>Mucor mucorales</i>	---	+_-	---	---	---
Enteric viruses					
Adenovirus	+_-	---	---	---	---
Norovirus	---	---	---	---	---
Rotavirus	---	---	---	---	---

Key: + Present; - Absent across the three month study period

Table 3: Cross-tabulation of relationship between Viruses, *Escherichia coli* and Protozoans

	<i>Escherichia coli</i>		<i>Cryptosporidium parvum</i>		<i>Giardia lamblia</i>	
	Present	Absent	Present	Absent	Present	Absent
Rotavirus	Present	0	0	0	0	0
	Absent	6	9	0	15	0
Adenovirus	Present	0	1	0	1	0
	Absent	6	8	0	14	0
Norovirus	Present	0	0	0	0	0
	Absent	6	9	0	15	0




Fig. 1: Percentage of bacteria isolated from sachet water




Fig. 2. Percentage of fungi isolated from packaged sachet water brands

REFERENCES

- Adekunle, L., Sridhar, M., Ajayi, A., Oluwade, P. and Olawuyi, J. (2004). An assessment of the health and social implications of sachet water in Ibadan, Nigeria: A public health challenge. *African Journal of Biomedical Research* 7:5-8.
- Armon R.H. (2015). *Indicators of waterborne viruses*. In: Armon R., Hänninen O. (eds) Environmental Indicators. Springer, Dordrecht.
- Arnold, B.F. and Colford, J.M. (2007). Treating water with chlorine at point-of-use to improve water quality and reduce child diarrhoea in developing countries: a systematic review and meta-analysis. *American Journal of Tropical Hygiene* 76:354-364.
- Arowolo, K.O., Ayolabi, C.I., Lapinski, B., Santos, J.S., Raboni, S.M. (2019). Epidemiology of enteric viruses in children with gastroenteritis in Ogun State, Nigeria. *Journal of Medical Virology* 91(6): 1022-1029.
- Ashbolt, N.J. (2015). Microbial contamination of drinking water and human health from community water systems. *Curr Environ Health Rep.* 2(1):95-106.
- Barnett, H.L. and Hunter, B.B. (2006). Illustrated genera of imperfect fungi. 4th Edition, The American Phytopathological Society, St. Paul Minnesota.
- Bello, R; and Falano, O.C. (2017). Interpretation of aeromagnetic anomalies over Abeokuta, Southwest Nigeria, using spectral depth technique. *J. Appl. Sci. Environ. Manage.* 21 (2) 218-222.
- Bernstein, D.I. (2009). Rotavirus overview. *The Pediatric Infectious Disease Journal* 28:50 -53.
- Blanton, L.H., Adams, S.M., Beard, R.S., Wei, G.E., Bulens, S.N., Widdowson, M.A., Glass, R.I. and Monroe S.S.(2006). Molecular and epidemiologic trends of caliciviruses associated with outbreaks of acute gastroenteritis in the United States, 2000-2004. *Journal of Infectious Diseases* 193:413-421.
- Boschi-Pinto, C., Velebit, L. and Shibuya, K. (2008). Estimating child mortality due to diarrhea in developing countries. *Bulletin World Health Organization* 86:710 - 717.
- Chinyelu, A., Samuel, O., Chinyere, N., Nwora, A. and Christine, I. (2010). Parasites associated with sachet drinking water (pure water) in Awka, South-Eastern, Nigeria. *Sierra Leone Journal of Biomedical Research* 2:(1) 23-27.
- Cho, H.B., Lee, S.H., Cho, J.C. and Kim, S.J. (2000). Detection of adenoviruses and enteroviruses in tap water and river water by reverse transcription multiplex PCR. *Can. J. Microbiol.* 46:417-424.
- Dada, A.C. (2009). Sachet water phenomenon in Nigeria: Assessment of the potential health impacts. *Afr. J. Microbiol. Res.* 3(1):015-021.
- Ezeugwunne, I.P., Agbakoba, N.R., Nnamah, N.K. and Anahalu, I.C. (2009). Prevalence of bacteria in packaged sachets water sold in Nnewi, South East Nigeria. *World Journal of Dairy and Food Sciences* 4:19-21.
- Fong, T.T. and Lipp, E.K. (2005). Enteric viruses of humans and animals in aquatic environments: Health risks, detection, and potential water quality assessment tools. *Microbiol. Molec. Biol. Rev.* 69:357-371.
- Fongaro, G., do Nascimento, M., Rigotto, C., Ritterbusch, G., da Silva, A.D., Esteves, P.A. and Barardi C.R.M. (2013). Evaluation and molecular characterization of human adenovirus in drinking water supplies: viral integrity and viability assays. *Virology Journal* 10:166.
- Garrity, G.M., Bell, J.A. and Lilburn T. 2004. Taxonomic outline of the prokaryotes. *Bergey's Manual® of*

- Systematic Bacteriology. Second edition. 399 pages.
- Genthe, B., Gericke, M., Bateman, B., Mjoli, N., Kfir, R. (1995). Detection of enteric adenoviruses in South-African waters using gene probes. *Water Sci. Technol.* 31:345 - 350.
- Gibson-Schwab, K., Opryszko, M., Schissler, J. and Guo Y. (2011). Evaluation of human enteric viruses in surface water and drinking water resources in Southern Ghana. *American Journal of Tropical Hygiene* 84(1) 20-29.
- Gotfred-Rasmussen, H., Lund, M., Enemark, H.L. Erlandsen, M. and Petersen, E. (2016). Comparison of sensitivity and specificity of 4 methods for detection of *Giardia duodenalis* in feces: immunofluorescence and PCR are superior to microscopy of concentrated iodine-stained samples. *Diagnostic Microbiology and Infectious Disease* 84 (3):187-190.
- Goettlich, E., van der Lubbe, W., Lange, B., Fiedler, S., Melchert, I., Reifenrath, M., Flemming, H.C., de Hoog, S. (2002). Fungal flora in groundwater-derived public drinking water. *International Journal of Hygiene and Environmental Health* 205: 269–279.
- Green, K.Y. (2007). *Caliciviridae: The Noroviruses*. In: Knipe DM, Howley PM (eds.). *Fields Virology*, 5th Edition. Lippincott-Williams & Wilkins Publishers, Philadelphia p949-979.
- Hageskal, G., Lima, N., and Skaar, I. (2009). The study of fungi in drinking water. *Mycological Research* 113 (2):165-172.
- Hssaine, A., Gharbi, J., Harrath, R., Harrak, R., Chait, A., Aouni, M and Hafid J. (2011). In search of enteroviruses in water media in Marrakech, Morocco. *African Journal of Microbiological Research* 5(16): 2380-2384.
- Jack, S., Bell, D. and Hewitt, J. (2013). Norovirus contamination of a drinking water supply at a hotel resort. *New Zealand Medical Journal* 126:98-107.
- Jiang, S.C. (2006). Human adenoviruses in water: occurrence and health implications: a critical review. *Environ. Sci. Technol.* 40(23):7132-40.
- Jonathan, S.G., Oghodero, O. and Asemoloye, M.D. (2016). Incidence of moulds in treated and untreated drinking water of selected Local Governments in Ibadan, South-Western Nigeria. *Res. Rev. Biosci.* 11(3):108.
- Kageyama, T., Kojima, S., Shinohara, M., Uchida, S., Fukushi, S., Hoshino, F. et al. (2003). Broadly reactive and highly sensitive assay for Norwalk-like viruses based on real-time quantitative reverse transcription-PCR. *J. Clin. Microbiol.* 41:1548-1557.
- Kapikian, A.Z., Hoshino, Y. and Chanok, R.M. (2001). *Rotaviruses*. In: *Fields Virology*, 4th edn, Edited by DM. Knipe & PM. Howley. Philadelphia, PA: Lippincott Williams & Wilkins. Pp1787-1833.
- Karim, M.R., Pontius, F.W. and LeChevallier M.W. (2004). Detection of Noroviruses in Water – Summary of an International Workshop in Water. *Journal of Infectious Diseases* 189:21-28.
- Kwakyenuako, G., Borketey, P.B., Mensah-Attipee, I., Asmah, R.H. and Ayeh-Kumi, P.F. (2007). Sachet drinking water in Accra: The Potential Threats of Transmission of Enteric Pathogenic Protozoan Organisms. *Ghana Medical Journal* 41:62-67.
- LeChevallier, M., William, D. and Ramon, G. (1991). *Giardia* and *Cryptosporidium* species in filtered drinking water supplies. *App. and Environ. Microbiol.* 57 (9):2617-2621.
- Matthijnsens, J., Ciarlet, M., Rahman, M., Attoui, H., Bányai, K., Estes, M.K.,

- Gentsch, J.R. *et al.* (2008). Recommendations for the classification of group rotaviruses using all 11 genomic RNA segments. *Archives of Virology* 153:1621-1629.
- Maunula, L., Miettinen, I.T. and von Bonsdorff C-H. (2005). Norovirus outbreaks from drinking water. *Emerging Infectious Diseases* 11(11):1716-21.
- Maunula, L. (2007). Waterborne norovirus outbreaks. *Future Virol.* 2(1):101-112.
- Mbah, C.E., Muhammed, H. (2015). Examination of two brands of sachet water and tap water for pathogenic microorganisms. *World Rural Observ.* 7(1):1-7.
- Mena, K.D. and Gerba, C.P. (2009). Waterborne adenovirus. *Reviews of Environmental Contamination and Toxicology* 198:133-167.
- Mhlongo, N.T., Tekere, M., and Sibanda, T. (2019). Prevalence and public health implications of mycotoxigenic fungi in treated drinking water systems. *Journal of Water and Health* 17(4): 517-531.
- Miagostovich, M., Ferreira, F., Guimaraes, F., Fumian, T., Diniz-Mendes, L., Luz, S. *et al.* (2008). Molecular detection and characterization of gastroenteritis viruses occurring naturally in the stream waters of Manaus, Central Amazônia, Brazil. *App. Journal of Applied Microbiology and Environmental Microbiology* 7(2):375-382.
- National Population Commission (NPC) (2006). Nigeria National Census: Population Distribution by Sex, State, LGAs and Senatorial District: 2006 Census Priority Tables (Vol. 3). <http://www.population.gov.ng/index.php/publication/140-popn-distri-by-sex-state-jgas-and-senatorial-distr-2006>.
- Novak Babič, M., Gunde-Cimerman, N., Vargha, M., Tischner, Z., Magyar, D., Veríssimo, C., Sabino, R., Viegas, C., Meyer, W., and Brandão, J. (2017). Fungal contaminants in drinking water regulation? A tale of ecology, exposure, purification and clinical relevance. *International Journal of Environmental Research and Public Health* 14(6), 36.
- Novak Babič, M., Zupančič, J., Brandão, J., and Gunde-Cimerman, N. (2018). Opportunistic water-borne human pathogenic filamentous fungi unreported from food. *Microorganisms* 6, 79.
- Odeyemi, O.A. (2015). Bacteriological safety of packaged drinking water sold in Nigeria: public health implications. Springerplus. 4:642.
- Odjegba, E.E.; Idowu, O.A, Ikenweibe, N.B., Martins, O., Sadeeq, A.Y. (2015). Public Perception of Potable Water Supply in Abeokuta South west, Nigeria. *J. Appl. Sci. Environ. Manage.* 19 (1):1-9.
- Oladipo, I.C., Onyenika, I.C. and Adebisi, A.O. (2009). Microbial analysis of some vended sachet water in Ogbomoso, Nigeria. *African Journal of Food Science* 3(12):406 - 412.
- Oliveira, H. M. B., Santos, C., Paterson, R. R. M., Gusmão, N. B., and Lima, N. (2016). Fungi from a groundwater-fed drinking water supply system in Brazil. *International Journal of Environmental Research and Public Health* 13(3).
- Olowofela, J. A., Akinyemi, O.D., Idowu, O.A., Olurin, O.T. and Ganiyu, S.A. (2013). Estimation of magnetic basement depth beneath the Abeokuta Area, South West Nigeria, using aeromagnetic data. *Asian Journal of Earth Sciences* 5(3):70-78.
- Paterson, R.R.M. and Lima, N. 2015. *Molecular biology of food and water borne mycotoxigenic and mycotic fungi of humans*. Taylor and Francis Group, CRC Press: Baton Rouge, LA, USA. p.618.

- Paterson, R.R.M. (2019). Editorial for the special issue: Human pathogenic filamentous fungi from food/water and mycotoxin from water. *Microorganisms* 7:21.
- Pusch, D., Oh, D., Wolf, S., Dumke, R., Schroter-Bobsin, U., Höhne, M., Rösk, I. and Schreier, E. (2005). Detection of enteric viruses and bacteria indicators in German environmental waters. *Archive of Virology* 150: 929-947.
- Reynolds, K., Mena, K. and Gerba, C. (2008). Risk of waterborne illness via drinking water in the United States. *Reviews of Environmental Contamination and Toxicology* 192: 117-158.
- Seitz, S.R., Leon, J.S., Schwab, K.J., Lyon, G.M., Dowd, M., McDaniels, M., Abdulhafid, G., Fernandez, M.L., Lindesmith, L.C., Baric, R.S., and Moe, C.L. (2011). Norovirus infectivity in humans and persistence in water. *Applied and Environmental Microbiology* 77(19):6884-6888.
- Shittu, O.B., Adeniran, S.A., Afolabi, O.R. and Sam-Wobo. S.O. (2013). Random Amplified Polymorphic DNA typing of multidrug-resistant clinical and environmental *Pseudomonas aeruginosa* strains from South-West, Nigeria. *Journal of Natural Sciences, Engineering and Technology* 12:13-24.
- Shittu, O.B., Adeniran, S.A., Afolabi, O.R. and Sam-Wobo. S.O. (2014). Risk surveillance of multidrug resistant *Pseudomonas aeruginosa* in water and plasmid relatedness with clinical strains in Abeokuta, South West Nigeria. *Journal of Natural Sciences, Engineering and Technology* 13:44 - 57.
- Shittu, O.B., Adelaja, O.M., Obuotor, T.M., Sam-Wobo, S.O. and Adenaike, A.S. (2016) PCR-Internal Transcribed Spacer (ITS) genes sequencing and phylogenetic analysis of clinical and environmental *Aspergillus* species associated with HIV-TB coinfecting patients in an hospital in Abeokuta, southwestern Nigeria. *African Health Sciences* 16(1): 141 - 148.
- Siqueira, V.M., Oliveira, H.M.B., Santos, C., Paterson, R.R.M., Gusmão, N.B. and Lima, N. (2011). Filamentous fungi in drinking water, particularly in relation to biofilm formation. *International Journal of Environmental Research and Public Health* 8(2): 456-469.
- Swenson, P., Wadell, G., Allardm, A. and Hierholzer, J. (2003). *Adenoviruses*. In: Yolken RH, Landry ML, Smith TF, Waner JL. (ed.). *Manual of Clinical Microbiology*, Vol. II, 8th ed. ASM Press, Washington, DC. pp. 1404-1417.
- Szewzyk U, Szewzyk R, Manz W, Schleifer KH. (2000). Microbiological safety of drinking water. *Annu Rev Microbiol.* 54:81-127.
- Teunis, P.F., Moe, C.L., Liu, P., Miller, S.E., Lindesmith, L., Baric, R.S., Le Pendu, J., and Calderon. R.L. (2008). Norwalk virus: how infectious is it? *J. Med. Virol.* 80(8):1468-76.
- Thliza, I.A., Khan, A.U. and Dangora, D.B. (2015). Fungi Contamination of Some Selected Brands of Sachet Water Marketed in Ahmadu Bello University, Zaria, Nigeria. *Journal of Microbiology Research* 5(1): 23-30.
- United Nations International Child Educational Fund (UNICEF). (2008). *The State of the World's Children: Child Survival 2008*. Available at: <http://www.unicef.org/sowc08/>.
- United States Environmental Protection Agency (USEPA) (1998). *Drinking water contamination candidate list. Federal Regulations* 63:10274 - 10287.
- United States Environmental Protection Agency (USEPA). (2005). *Drinking water contaminant candidate list 2; final notice. Fed. Regist.* 70:9071-9077. <http://www.epa.gov>

- /fedrgstr/EPA
WATER/2005/February/Day-
24/w3527. htm.
- United States Environmental Protection Agency (USEPA). (2009). *United States Environmental Protection Agency Contaminant Candidate List 3*.
http://water.epa.gov/scitech/drinking_water/dws/ccl/ccl3.cfm.
- Upfold, N.S., Luke, G.A. and Knox, C. (2021). Occurrence of human enteric viruses in water sources and shellfish: a focus on Africa. *Food Environ Virol.* 13: 1 - 31.
- Van Heerden, J., Ehlers, M.M., Van Zyl, W.B. and Grabow, W. (2003). Incidence of adenoviruses in raw and treated water. *Water Research* 37: 3704-3708.
- Van Heerden, J., Ehlers, M.M., Van Zyl, W.B. and Grabow, W. (2004). Prevalence of human adenoviruses in raw and treated water. *Water Science Techn.* 50: 39-43.
- Van Heerden, J., Ehlers, M.M., Van Zyl, W.B. and Grabow, W. (2005). Detection and risk assessment of adenoviruses in swimming pool water. *Journal of Applied Microbiology* 99: 1256-1264.
- Verheyen, J., Timmen-Wego, M., Laudien, R., Boussaad, I., Sen, S., Koc, A., Uesbeck, A., Mazou, F. and Pfister, H. (2009). Detection of adenoviruses and rotaviruses in drinking water sources used in rural areas of Benin, West Africa. *Applied Environ. Virol.* 75:2798-2801.
- World Health Organization (WHO). (2003). *Guidelines for environmental surveillance of poliovirus circulation. Vaccines and Biologicals.* World Health Organization, Geneva, Switzerland. Pp 1- 19.
- World Health Organization (WHO). (2008). *Guidelines for drinking water quality.* World Health Organization, Geneva, Switzerland: Pp. 1- 668.
- World Health Organization and United Nations International Child Educational Fund (WHO/UNICEF). (2006). *Meeting the MDG drinking water sanitation target: the urban and rural challenges of the decade.* Available at http://www.who.int/water_sanitation_health/monitoring/jmp2006/en/index.html.
- Xagorarakis, I., Kuo, D.H., Wong, K, Wong, M and Rose, J.B. (2007). Occurrence of human adenoviruses at two recreational beaches of the Great Lakes. *Appl. Environ. Microbiol.* 73:7874-7881.
- Yu, J.R., Lee, S.U., and Park, W.Y. (2009). Comparative sensitivity of PCR primer sets for detection of *Cryptosporidium parvum*. *The Korean Journal of Parasitology* 47(3): 293-297.