

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/338422546>

Alteration in Hormonal level and Testicular Histomorphology in Rats Treated with Alcohol (ALC) and Cannabinol (CBN)

Article · December 2019

CITATIONS

0

READS

115

6 authors, including:



Temitayo O. Ajayi

University of Ibadan

23 PUBLICATIONS 139 CITATIONS

SEE PROFILE



Charles Oluwafemi Faeji

Afe Babalola University

19 PUBLICATIONS 141 CITATIONS

SEE PROFILE



Akele Richard

Afe Babalola University

67 PUBLICATIONS 281 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Water Management [View project](#)



Drinking water quality [View project](#)

UNIVERSITY OF IBADAN LIBRARY

Alteration in Hormonal level and Testicular Histomorphology in Rats Treated with Alcohol (ALC) and Cannabinol (CBN)

Akintayo C.O^{1*}, Ajayi T.O², Akinsomisoye S.O³, Faeji C.O⁴,
Adewumi A.F⁵, Akele Y.R⁵.

^{1*}Department of Physiology, College of Medicine and Health Sciences, Afe Babalola University Ado Ekiti, Ekiti State, Nigeria.

²Department of Pharmacognosy, Faculty of Pharmacy, University of Ibadan. Nigeria.

³Department of Physiological Sciences, Obafemi Awolowo University, Ile Ife, Osun State, Nigeria.

⁴Department of Medical Microbiology and Parasitology, College of Medicine and Health Sciences, Afe Babalola University, Ado Ekiti, Ekiti State, Nigeria

⁵Department of Medical Laboratory Science, College of Medicine and Health Sciences, Afe Babalola University, Ado Ekiti, Ekiti State, Nigeria

^{1*}**Corresponding author:**

Akintayo Christopher.

Email: tobajoe2001@yahoo.com

Published: 27 December 2019

Copyright © Akintayo et al.

Cite this article: Akintayo C.O., Ajayi T.O., Akinsomisoye S.O., Faeji C.O., Adewumi A.F. & Akele Y.R. (2019). Alteration in Hormonal level and Testicular Histomorphology in Rats Treated with Alcohol (ALC) and Cannabinol (CBN). *European International Journal of Science and Technology*, 8(11), 41-49.

ABSTRACT

This study explores the effects of chronic alcohol and cannabinol exposure on the testosterone levels and the histomorphological integrity/cytoarchitecture of the rats' testes. Adults male Wistar rats were randomly divided into control (distilled water), methanol (2mg/kg bw), alcohol (3g/kg bw), cannabinol (10mg/kg bw) and alcohol (3g/kg bw) plus cannabinol (10mg/kg bw) groups respectively. Alcohol and cannabinol were administered orally twice daily for 52 days (spermatogenic cycle in rats) and at the end of treatment, male reproductive organ (testis) was removed and cleared of adherent tissue and then fixed for histological examination. Blood samples were collected via retro-orbital sinus for hormone (testosterone) assay. Serum testosterone levels were measured using the enzyme-immunoassay (E.I.A.) technique. Hormonal assay showed significant reductions in the levels of testosterone (T) ($p < 0.05$) in the alcohol alone, cannabinol alone and in the alcohol plus cannabinol treated groups. The Histological analysis of the treated groups showed severe reduction of the spermatogenic cells. The present study showed that following chronic alcohol and/or cannabinol administration the results showed significant reduction ($p < 0.05$) in testosterone levels and a detrimental effect on the histomorphology of the testes. Alcohol and/or cannabinol therefore exhibit inhibitory effects causing inhibition of testosterone as observed in this study.

Keywords: Alcohol, cannabinol, spermatogenesis, hormone, histomorphology.

1. Introduction

The use of drugs is on the increase among people of reproductive age. Alcohol and cannabinoids belong to the commonly abuse drugs among the youths and belongs to the most widely used drugs which can suppress reproductive functions and sexual behavior both in human and in experimental animals (Abel, 1980; Talabi *et al.*, 2011). Alcohol consumption is often associated with poor semen production and sperm quality (Symons *et al.*, 1976; Mendelson *et al.*, 1977). Chronic consumption of alcohol can result to fertility disturbances (low sperm count, sperm motility and reduced serum testosterone level, testicular atrophy) both in men and laboratory animals (Abel, 1980; Mendelson *et al.*, 1977). However, cannabinol also belongs to the class of illicit drug which can evoke deleterious effects on reproductive activities both in males and in females. Researchers have investigated that tetrahydro-cannabinol induced blockage of the release of gonadotropins and have resulted in lowered LH and FSH, both of which are responsible for the reduction of testosterone production by the Leydig cells of the testis (Dalterio and Bartke, 1981). Testosterone is essential in males during spermatogenesis, as it is being secreted by the testicular Leydig cells under the influence of luteinizing hormone (LH) and acts on the seminiferous epithelium; thus enhances spermatogenesis. The Leydig cells produce testosterone which is essential for the development of the male reproductive system. Interference with Leydig cell testosterone production in male adults causes diminution in male sexual behavior, atrophy of the testis, epididymis and accessory sex organs and impairment of spermatogenesis. However, the role of testosterone on spermatogenesis could be suppressed as a result of exposure to illicit drugs, recreational drugs and other synthetic drugs. Various substances have been implicated as Leydig cell toxicants and some of these substances act directly on Leydig cells, while others could act directly on the hypothalamic-pituitary-testicular axis. Acute and chronic administration of cannabinoids including tetrahydrocannabinol (THC), cannabinol (CBN), and cannabidiol (CBD) has been reported to decrease sexual functions both in male and female rodents as well as in primates. The effects of cannabinoids on both male and female reproductive system have been studied (Marks, 1973; Symons *et al.*, 1976; Kolodny *et al.*, 1976). However, study has not

revealed the combined administration of alcohol and cannabinal on reproductive hormone levels and the testicular cytoarchitecture in animal model. This study seeks to evaluate testosterone level and testicular histomorphology in rats exposed to alcohol and cannabinal.

2. MATERIALS AND METHODS

2.1 Experimental Design

2.1.1 Animals

Thirty (30) healthy male Sprague-Dawley rats weighing 180-200 g (12 weeks old) were randomly selected from the Laboratory Animal Center of NIPRD, Abuja. The rats were fed on standard diet (Ladokun Feed and Flour Mills Ltd), water was given ad libitum and maintained under standard conditions. The animal room was well ventilated with a temperature range of 25 - 27°C under light/dark 12-12 h photoperiodicity. The rats were grouped into 5 groups of 5 rats each. The use of the animals was approved by the Experimental Ethic Committee on Animal Use of National Institute of Pharmaceutical and Research Development (NIPRD), Abuja, Nigeria and was conducted in accordance with the guidelines of the Declaration of Helsinki.

2.2 Experimental Groups

Group 1 was the control (distilled water), group 2 rats received (2mg/kg bw), group 3 was the alcohol treated group (3g/kg bw), group 4 was cannabinal treated group (10mg/kg bw), and group 5 was treated with Alcohol (3g/kg bw) plus cannabinal (10mg/kg bw). Drugs and vehicle administration was done orally using an oral cannula for 48 days. Drugs given to each rat was based on its body weight (in grams). The animals were weighed before and after the treatment period. The drug treated groups were sacrificed after the 52 days of administration.

2.2.1 Anesthetic protocol and autopsy

The rats were examined on a daily basis throughout the experimental period for signs of toxicity. At the end of the experiment the rats were sacrificed via cervical dislocation under 25% urethane anesthesia (0.6 ml/ 100 g bw).

2.2.2 Organ Collection

The animals were sacrificed by cervical dislocation and were dissected and reproductive organ of interest (testis) was removed, cleared of adherent tissue and weighed immediately using electronic weighing balance.

2.2.3 Hormone assay

Blood was collected from each rat via cardiac puncture from which the serum was separated. Serum testosterone hormone levels were measured using the enzyme-immunoassay (E.I.A.) technique. The E.I.A. kits were produced by inteco (London, UK) and the optical density was read using a spectrophotometer (Jenway, 6300 Spectrophotometer, UK) that was sensitive at wavelengths between 492–550 nm.

2.2.4. Histomorphological Evaluation

The testis from each group was harvested and fixed in bouin's fluid for six hours and then transferred into 10% buffered formalin for 24 hours after which it was transferred to 70% alcohol for dehydration. The tissues were passed through 90% and absolute alcohol and xylene for

different durations before they were transferred into two changes of molten paraffin wax for 1 hour each in an oven at 65 o C for infiltration. They were subsequently embedded and serial sections cut using rotary microtome at 5 microns. The tissues were picked up with albumenised slides and allowed to dry on hot plate for 2 min. The slides were dewaxed with xylene and passed through absolute alcohol (2 changes); 70% alcohol, 50% alcohol and then to water for 5 min. The slides were then stained with haematoxylin and eosin. The slides were mounted and the photomicrographs of the slides were then taken at magnifications of $\times 100$.

2.2.5. Statistical Analysis

Data analysis was carried out using one way analysis of variance (ANOVA) supported by Newman-Keuls test when followed by Turkey's multiple comparison pair with comparison was done between groups. Results were presented as mean \pm SEM, and the differences were considered significant at $p < 0.05$.

3. Results




Figure 1. Testosterone concentrations of rats exposure to alcohol and/or cannabinol

3.1.1 Histomorphology of experimental animals




Figure 1: Photomicrography of a testicular section treated with alcohol and/or cannabiniol in Wistar rats

A is designated as the control group showing normal seminiferous tubules (white arrows) with full maturation of spermatocytes. The germinal cell layer (spanned) shows developmental stages. The interstitial cells appear normal (slender arrow). B, is the group that served as the vehicle (methanol) for cannabiniol, the testicular histology showed normal cytoarchitecture with normal seminiferous tubules and both primary and secondary spermatocytes. C, D and E were treated with alcohol, cannabiniol and alcohol+cannabiniol respectively; the cytoarchitecture showed very few seminiferous tubules showing incomplete maturation with widening of the interstitial space with interstitial cells showing hyperplasia (black arrow). (X100, H&E).

4. Discussion

Substances of abuse exhibit toxic effects in the reproductive functions in both humans and laboratory animals. Cannabinol and alcohol exerted dramatic reproductive toxicity by lowering the testosterone levels and causing a deleterious effect on the testes. Testosterone is the primary androgen that controls the functional activity of virtually all male reproductive tract structures. Spermatogenesis in seminiferous tubules, sperm maturation in the epididymis and the secretory activity of the accessory sex glands all require adequate levels of testosterone, although spermatogenesis in the rat can be maintained with intra-testicular testosterone concentrations as low as 20% of normal values. Oral administration of Δ^9 -tetrahydrocannabinol, the major active ingredient of cannabis (Marijuana), increases both plasma testosterone and LH concentrations in male mice as reported by (Daltero *et al.*, 1982, 1984). Testicular histomorphological examination of groups treated with alcohol and/or cannabinol showed degeneration of spermatogenic cells in the seminiferous tubules and in complete spermatogenesis. However, there was an improvement in the testicular architecture in rats co-treated with quercetin alongside alcohol and/or cannabinol. The alteration in reproductive functions may also be linked to the decrease in the mass of the testes and inhibition of testicular androgenesis (testosterone), likewise reducing the serum concentrations of gonadotrophins (FSH and LH) as revealed in the present study. Testosterone is essential for the sequential stages of spermatogenesis and also responsible in the maintenance of spermatogenesis; it aids in the growth of external genitalia (penis, scrotum and other accessory sex organs including the genital ducts, seminal vesicles and prostate gland respectively). Moreover, the reduction in serum testosterone levels in the alcohol and/or nicotine treated rats could be associated with the inhibitory effect on the pituitary gland thereby altering reproductive functions. Although, the regulation of testosterone production or secretion is regulated by Luteinizing hormone (LH) or interstitial cell stimulating hormone (ICSH). LH is important in stimulating the Leydig cells and the quantity of testosterone is proportional to the concentration of LH. However, the reduction in the serum testosterone concentration observed in the rats treated with alcohol and/or cannabinol could be linked to the reduction in the serum levels of LH. The effect appears to be a direct response of the glandular tissues involved, as the release of testosterone does not require the presence of the pituitary, and the administration of testosterone does not inhibit the release of LH. The effect on testosterone is biphasic, causing rapid sustained increases in plasma levels at low doses and subsequent decreases at higher doses. Daltero *et al.*, 1984, suggested that this may be responsible for the reported aphrodisiac effects of cannabis, which are arousal at low doses or in occasional users, followed by suppression of libido at high concentrations. The present result is in agreement with the study by Purohit *et al.*, (1979, 1980) showed that both tetrahydrocannabinol and cannabinol decreased ventral prostate, seminal vesicular weight and plasma concentrations of testosterone and dihydrotestosterone in adult male rats after 10 days of subcutaneous injection at a dose of 10ng/kg of body weight. The serum testosterone levels of rats' treatment with cannabinol at a dose of 10mg/kg of body weight showed a marked significant reduction in comparison to the control counterpart. Studies by Marks, 1973; Kolodny *et al.*, 1976, also revealed that exposure to cannabinoids involving a mice model could be confusing because cannabinoids have been shown to both increase and decrease serum testosterone concentrations in peripheral blood depending on the dosage, time of blood sampling, route of drug administration and the interactions with the gonadotropins (FSH and LH) (Dalterio *et al.*, 1976, 1979; Dalterio and Bartke, 1981). Reports by Dalterio *et al.*, 1979; Frias *et al.*, 2000, revealed alterations in the function of the hypothalamo-hypophyseal-gonadal axis in adult male mice following prenatal exposure to cannabinoids. This study is in alignment with earlier reports revealing the detrimental effects of

alcohol and cannabinol administration on the reproductive activities evident by the alteration derangement in the cytoarchitecture of the epididymis and testis respectively (Akintayo *et al.*, 2015).

4.1 Conclusion

In conclusion the present study demonstrates that following chronic administration of alcohol and /or cannabinol caused reduction in serum testosterone levels and deleterious effects on the cytoarchitecture of the rats' testes. Thus, alcohol and cannabinol are testicular toxins and are believed to produce site-specific lesion at such sites as the Leydig cell, germ cells and as well as Sertoli cell.

References

- [1] Abel, E.L. (1980). A review of alcohol's effects on sex and reproduction. *Drug Alcohol Depend*; 5:321–332.
- [2] Akintayo, C.O., Karga, S.V., Ayodele, M.J., Sanya, J.O., Adekeye, A.O. and Ibitayo, A.O. (2015). Histomorphological Examination of Reproductive and Accessory Organs Exposed to Alcohol and Cannabinol in Male Adult Rats. *European International Journal of Science and Technology* 4 (5); 47-55.
- [3] Burstein, S., Hunter, S.A., Shoupe, T.S. (1979). Site of Inhibition of Leydig cell testosterone synthesis by Δ^1 -Tetrahydrocannabinol. *Mol. Pharmacol*; 15: 633-640.
- [4] Dalterio, S., Bartke, A., Burstein, S. (1976). Cannabinoids inhibit testosterone secretion by mouse testes in vitro. *Science*; 196: 1472-1473.
- [5] Dalterio, S. and Bartke, A. (1979). Perinatal exposure to cannabinoids alters male reproductive function in mice. *Science*; 205:1420-1422
- [6] Dalterio, S., Bartke, A., Mayfield, D. (1981). Δ^9 -Tetrahydrocannabinol increases plasma concentrations in mice. *Science*; 213: 581-582.
- [7] Dalterio, S., Badr, F., Bartke, A., Mayfield, D. (1982). Cannabinoids in male mice: effects on fertility and spermatogenesis. *Science*; 216: 315-316.
- [8] Dalterio, S., Badr, F., Mayfield, A. D., Bartke, A. (1984). Early Cannabinoid exposure influences neuroendocrine and reproductive functions in male mice: prenatal exposure. *Pharmacol Biochem Behav*; 20: 107-133.
- [9] Frias, J., Rodriguez, R., Torres, J.M., Ruiz, E., Ortega, E. (2000). Effects of acute alcohol intoxication on pituitary gonadal axis hormones, pituitary adrenal axis hormones, beta-endorphin and prolactin in human adolescents of both sexes. *Life Sci*; 67: 1081–1086.
- [10] Harclerode, J. (1984). Endocrine effects of Marijuana in the male: preclinical studies. *NIDA Res. Monogr*; 44: 46-64.

- [11] Jakubovic, A., McGeer, E.G., McGeer, P.L. (1978). Biochemical alterations in the Leydig cells of the rat testis in vitro: effects on testosterone and protein synthesis. In Marihuana:
- [12] Johns, A. (2001). Psychiatric effects of cannabis. *British Journal of Psychiatry*;178 :116 -122.
- [13] Biological Effects (Nahas, G.G. and Paton, W.D., Eds.), Pergamon Press, New York; 251-264.
- [14] Kim, J.H., Kim, H.J., Noh, H.S., Roh, G.S., Kang, S.S., Cho, G.J. *et al.*(2003). Suppression by ethanol of male reproductive activity. *Brain Res*; 989: 91–98.
- [15] Kolodny, R.C., Lessin, P., Torro, G., Masters, W.H., Cohen, S. (1976). Depression of plasma testosterone with acute marihuana administration. In: Braude M.C. Szara S, eds. *Pharmacology of Marihuana*. New York: Raven Press; 1: 217-225.
- [16] Lenzi, A., Lombardo, F., Gandini, L., Alfano, P., Dondero, F. (1993). Computer assisted sperm motility analysis at the moment of induced pregnancy during gonadotropin treatment for hypogonadotropic hypogonadism. *J Endocrinol Invest*; 16: 683–686.
- [17] List, A., Nazar, B., Nyquist, S., Harclerode, J. (1977). The effects of Δ^9 -tetrahydrocannabinol and caccabinol metabolism of gonadal steroids in the rats. *Drug Metab. Dispos*; 5: 268-272.
- [18] Marks, B.H. (1973). Δ^9 -tetrahydrocannabinol and luteinizing hormone secretion. *Prog Brain Res.*; 39: 331-338.
- [19] Martinez, M., Macera, S., de Assis, G.F., Pinheiro, P.F., Almeida, C.C. Tirapelli, L.F. *et al.* (2009). Structural evaluation of the effects of chronic ethanol ingestion on the testis of *Calomys callosus*. *Tissue Cell*; 41:199–205.
- [20] Mendelson, J.H., Mello, N.K., Ellingboe, J. (1977). Effects of acute alcohol intake on pituitary–gonadal hormones in normal human males. *J Pharmacol Exp Ther*; 202:676–682.
- [21] Meneguz, A., Fortuna, S., Lorenzini, P., and Volpe, M.T. (1999). Influence of urethane and ketamine on rat hepatic cytochrome P450 in vivo. *Exp Toxicol Pathol*; 51:392–396.
- [22] Midgley, A.R., Niswender, G.D. (1970). Radioimmunoassay of steroids. *Acta Endocrinol*; 147: 320-331.
- [23] Nagy, F., Pendergrass, P.B., Bowen, D.C., Yeager, J.C.A. (1986). Comparative study of cytological and physiological parameters of semen obtained from alcoholics and non-alcoholics. *Alcohol Alcohol*; 21: 17–23.
- [24] Purohit, V., Singh, H.H., Ahluwalia, B.S. (1979). Evidence that the effects of methadone and marihuana on male reproductive organs are mediated at different sites in rats. *Biol Repro*; 20:1039-1044.

- [25] Purohit, V., Ahluwalia, B.S., Vigersky, R. (1980). Cannabinoids inhibits dihydrotestosterone binding to androgen receptors in the rat ventral prostate. *Endocrinology*; 197: 848-850.
- [26] Rouach, H., Fatacciolo, V., Gentil, M., French, S.W., Morimoto, M., Nordmann, R. (1997). Effect of chronic ethanol feeding on lipid peroxidation and protein oxidation in relation to liver pathology. *Hepatology*; 25: 351–355.
- [27] Selvage, D., Hales, D., Rivier, C. (2004). Comparison between the influence of the systemic and central injection of alcohol on leydig cell activity. *Alcohol Clin Exp Res*; 28: 480–488.
- [28] Sierksma, A., Sarkola, T., Eriksson, C., van der Gaag, M., Grobbee, D., Hendriks, H. (2004). Effect of moderate alcohol consumption on plasma dehydroepiandrosterone sulfate, testosterone, and estradiol levels in middle-aged men and postmenopausal women: a diet-controlled intervention study. *Alcohol Clin Exp Res*; 28: 780–785.
- [29] Silva, S.M., Santos-Marques, M.J., Madeira, M.D. (2009). Sexually dimorphic response of the hypothalamic–pituitary–adrenal axis to chronic alcohol consumption and withdrawal. *Brain Res*; 1303: 61–73.
- [30] Symons, A.M., Teale, J.D., Marks, V. (1976). Effects Δ^9 -tetrahydrocannabinol on the hypothalamic-pituitary gonadal system in the maturing rat. *J Endocrinol*; 68:43-44.
- [31] Talabi, A.R., Sarchesmeh, A.A., Khalili, M.A and Tabibreyad, N. (2011). Effects of ethanol consumption on chromatin condensation and DNA integrity of epididymal spermatozoa in rat. *Alcohol*; 45: 403–409.
- [32] Välimäki, M.J., Härkönen, M., Eriksson, C.J., Ylikahri, R.H. (1984). Sex hormones and adrenocortical steroids in men acutely intoxicated with ethanol. *Alcohol*;1: 89–93.
- [33] Vatsalya, V., Issa, J.E., Hommer, D.W., Ramchandani, V.A. (2012). Pharmacodynamic effects of intravenous alcohol on hepatic and gonadal hormones: influence of age and sex. *Alcohol Clin Exp Res*; 36: 207–213
- [34] Walter, M., Gerhard, U., Gerlach, M., Weijers, H.G., Boening, J., Wiesbeck, G.A. (2007). Controlled study on the combined effect of alcohol and tobacco smoking on testosterone in alcohol-dependent men. *Alcohol*; 42: 19–23.
- [35] Watanabe, K., Motoya, E., Matsuzawa, N., Funashashi, T., Kimura, T., Matsunaga, T., Arizozo, K., and Yamamoto, I. (2005). Marijuana extracts possess the effects like the endocrine disrupting chemicals. *Toxicology*; 206: 471-478.